

Convergence of functionals of sums of r.v.s to local times of fractional stable motions

P.Jeganathan
Indian Statistical Institute
Bangalore Center
8 th Mile, Mysore Road
Bangalore 560059
INDIA
e-mail: jegan@isibang.ac.in

First version: July, 2002.

This version: May, 2003.

Abbreviated title: Weak convergence to local times.

Summary. Consider a sequence $X_k = \sum_{j=0}^{\infty} c_j \xi_{k-j}$, $k \geq 1$, where c_j , $j \geq 0$, is a sequence of constants and ξ_j , $-\infty < j < \infty$, is a sequence of i.i.d. random variables belonging to the domain of attraction of a strictly stable law with index $0 < \alpha \leq 2$. Let $S_k = \sum_{j=1}^k X_j$. Under suitable conditions on the constants c_j it is known that for a suitable normalizing constant γ_n , the partial sum process $\gamma_n^{-1} S_{[nt]}$ converges in distribution to a linear fractional stable motion (indexed by α and H , $0 < H < 1$). A fractional ARIMA process with possibly heavy tailed innovations is a special case of the process X_k . In this paper it is established that the process $n^{-1} \beta_n \sum_{k=1}^{[nt]} f(\beta_n (\gamma_n^{-1} S_k + x))$ converges in distribution to $\left(\int_{-\infty}^{\infty} f(y) dy \right) L(t, -x)$, where $L(t, x)$ is the local time of the linear fractional stable motion, for a wide class of functions $f(y)$ that includes the indicator functions of bounded intervals of the real line. Here $\beta_n \rightarrow \infty$ such that $n^{-1} \beta_n \rightarrow 0$. The only further condition that is assumed on the distribution of ξ_1 is that either it satisfies the Cramér's condition or has a nonzero absolutely continuous component. The results have motivation in large sample inference for certain nonlinear time series models.

MSC 2000 Subject classifications. *Primary:* 60F05, 60G18, 60J55. *Secondary:* 62M10, 62J02.

Key words and phrases. Fractional stable motion, Fractional Brownian motion, Local time, Fractional ARIMA process, Functionals of sums of fractional ARIMA, Weak convergence to local times, Heavy tailed distributions.

1 Introduction

Consider a sequence ξ_j , $-\infty < j < \infty$, of i.i.d. random variables belonging to the domain of attraction of a strictly stable law with index $0 < \alpha \leq 2$. Define $X_k = \sum_{j=0}^{\infty} c_j \xi_{k-j}$, where $c_j, j \geq 0$, is a sequence of real numbers. Let $S_k = \sum_{j=1}^k X_j$, $k \geq 1$. Then under suitable conditions on the constants c_j it is known that for a suitable $0 < H < 1$ and for a suitable slowly varying function $u(n)$, the finite dimensional distributions of the process $(n^H u(n))^{-1} S_{[nt]}$ converge in distribution to those of a Linear Fractional Stable Motion (LFSM). See for example Kasahara and Maejima (1988). When $\alpha = 2$, the LFSM reduces to the Fractional Brownian Motion (FBM), and when $H = 1/\alpha$ it is taken to be the α -stable Lévy motion. (Definitions of the preceding processes are recalled in Section 2 below.)

Now for simplicity let $\gamma_n = n^H u(n)$. Let the constants β_n be such that $\beta_n \rightarrow \infty$ with $n^{-1} \beta_n \rightarrow 0$. In this paper it is established that the finite dimensional distributions of the process $n^{-1} \beta_n \sum_{k=1}^{[nt]} f(\beta_n (\gamma_n^{-1} S_k + x))$ (indexed by t and x) converge in distribution to those of $\left(\int_{-\infty}^{\infty} f(y) dy \right) L(t, -x)$, where $L(t, x)$ is the local time of the LFSM, for a wide class of functions $f(y)$ that in particular includes the indicator functions of bounded intervals. (The only further condition that will be assumed on the distribution of ξ_1 is that either it satisfies the Cramér's condition or has a nonzero absolutely continuous component.)

For the particular situation where the limit of $\gamma_n^{-1} S_{[nt]}$ is a Brownian motion or a FBM, some partial results in some form are available in Akonom (1993), and in Park and Phillips (1999, 2001) and Tyurin and Phillips (1999) where the motivation is an interesting development of a large sample theory in some time series models that have functions of the form $f(S_k)$ occurring as regressions. The present paper has the same motivation.

Major works for the i.i.d. situation $S_k = \sum_{j=1}^k \xi_j$ that are related to the approach of the present paper include Skorokhod and Slobodnjuk (1970) and Ibragimov (1984, 1985). The approach of Ibragimov (1984, 1985) (a thorough account of which is presented in the book by Borodin and Ibragimov (1995)) does not rely on the moment conditions, and hence implicitly on the restriction of the Brownian motion limit, of the earlier approach of Skorokhod and Slobodnjuk (1970) but mainly on the requirement of the attraction of $\sum_{j=1}^k \xi_j$ to stable laws. This approach is based on the representation

$$\frac{\beta_n}{n} \sum_{k=1}^n f(\beta_n \gamma_n^{-1} S_k) = \frac{1}{n} \sum_{k=1}^n f_n(\gamma_n^{-1} S_k) = \int_{-\infty}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n e^{-i u \gamma_n^{-1} S_k} \right) \hat{f}_n(u) du$$

where $f_n(y) = \beta_n f(\beta_n y)$ and $\hat{f}_n(u)$ is the Fourier transform of $f_n(y)$. The conditions sought (in the i.i.d. case) are naturally through the Fourier transform $\hat{f}_n(u)$ which in certain situations are then transformed in terms of $f_n(y)$ and/or on the distribution of ξ_1 .

The approach of the present paper involves the approximation that the difference

$$\frac{1}{n} \sum_{k=1}^n f_n(\gamma_n^{-1} S_k) - \frac{1}{n} \sum_{k=1}^n \int_{-\infty}^{\infty} f_n(\gamma_n^{-1} S_k + z\varepsilon) \phi(z) dz \rightarrow 0 \quad \text{in } \mathbf{L}^2 \quad (1)$$

as $n \rightarrow \infty$ first and then $\varepsilon \rightarrow 0$, where $\phi(z) = \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{z^2}{2}\right\}$. The approximating quantity $\frac{1}{n} \sum_{k=1}^n \int_{-\infty}^{\infty} f_n(\gamma_n^{-1} S_k + z\varepsilon) \phi(z) dz$ can be handled relatively easily. This approach has some advantages, especially for the situation of the present paper, for instance the required conditions for establishing the approximation (1) can be viewed directly through $f_n(y)$ and the distribution of ξ_1 themselves.

In Section 2 we recall the definition of the LFSM, state a result on the existence of its local time and also recall a result on the weak convergence of the process $\gamma_n^{-1} S_{[nt]}$. Section 3 contains the statements and the discussions of the main results, and Section 4 contains the proofs of them as consequences of (1). Section 5 contains auxiliary results for establishing the approximation (1). The proof of (1) constitutes Section 6.

Notations: The constants γ_n, β_n and the functions $f(y)$ and $f_n(y) = \beta_n f(\beta_n y)$ are exclusively used in the sense they are used above. Similarly, $\phi(z)$ in the sense of (1) above, b_n in the sense of (4) below, and $M_{h,\eta}(y)$ and $m_{h,\eta}(y)$ in the sense of (6) below. We let $\phi_\varepsilon(z) = \frac{1}{\varepsilon\sqrt{2\pi}} \exp\left\{-\frac{z^2}{2\varepsilon^2}\right\}$ (so that $\phi(z) = \phi_1(z)$). \widehat{K} stands for the Fourier transform of the measure K , $\mathbb{1}_A(\cdot)$ stands for the indicator function of the set A and \mathbb{R} stands for the real line. Convergence in \mathbf{L}^2 has the usual meaning of the convergence in mean-square. The notation C stands for a generic constant that may take different values even at different places of the same proof.

2 Preliminaries

Let $\{Z_\alpha(t), t \in \mathbb{R}\}$, $0 < \alpha \leq 2$ be a an α -stable Levy motion. This means $Z_\alpha(t)$ has stationary independent increments having a strictly α -stable distribution, that is, for $s < t$, $Z_\alpha(t) - Z_\alpha(s)$ has the characteristic function $\exp\left\{-(t-s)|u|^\alpha \left(1 + i\beta \operatorname{sign}(u) \tan\left(\frac{\pi\alpha}{2}\right)\right)\right\}$ where $|\beta| \leq 1$ with $\beta = 0$ when $\alpha = 1$. (Note that this definition of strict α -stability for the case $\alpha = 1$ differs from the usual one in that we take the shift parameter to be 0.) When $\alpha = 2$, $Z_\alpha(t)$ becomes the *Brownian Motion* with variance 2.

A process $\{\Lambda_{\alpha,H}(t), t \geq 0\}$ is called a *Linear Fractional Stable Motion* (LFSM) with Hurst parameter H , $0 < H < 1$, if it is given by

$$\Lambda_{\alpha,H}(t) = a \int_{-\infty}^0 \left\{ (t-u)^{H-1/\alpha} - (-u)^{H-1/\alpha} \right\} Z_\alpha(du) + a \int_0^t (t-u)^{H-1/\alpha} Z_\alpha(du)$$

where $Z_\alpha(t)$ is an α -stable Levy motion as above and a is a non-zero constant. When

$\alpha = 2$, the LFSM reduces to the *Fractional Brownian Motion* (FBM). See Samorodnitsky and Taqqu (1994) and Maejima (1989) for the details of LFSM.

We make the convention that in the case $H = 1/\alpha$ the LFSM $\{\Lambda_{\alpha,H}(t), t \geq 0\}$ is taken to be $\{Z_\alpha(t), t \geq 0\}$. It is important to note however that in this case the restriction $0 < H < 1$ is equivalent to that of $1 < \alpha \leq 2$.

Let $\{\zeta(t), t \geq 0\}$ be a real valued measurable process. Then a measurable process $\{L(t, x), t \geq 0, x \in \mathbb{R}\}$ is said to be a *local time* of $\{\zeta(t), t \geq 0\}$ if for each $t \geq 0$

$$\int_0^t \mathbb{I}_A(\zeta(s)) ds = \int_{-\infty}^{\infty} \mathbb{I}_A(x) L(t, x) dx \quad \text{for all Borel subset } A \text{ of } \mathbb{R} \quad (2)$$

with probability one. (Without loss of generality we take $\Lambda_{\alpha,H}(t)$ to be measurable.) For the symmetric LFSM the existence of $L(t, x)$ is known, see Kôno and Maejima (1991). For the general LFSM, we have the following result. (It may be noted that when $0 < \alpha \leq 1$ the local time for $\{Z_\alpha(t)$ does not exist; the case $1 < \alpha \leq 2$ is covered by the next result).

Theorem 0. *For a LFSM $\{\Lambda_{\alpha,H}(t), t \geq 0\}$ with $0 < \alpha \leq 2$ and $0 < H < 1$, there is a local time $L(t, x)$ such that for each t and x , $\lim_{\eta \downarrow 0} \frac{1}{\eta} \int_0^t \mathbb{I}_{[x, x+\eta)}(\Lambda_{\alpha,H}(s)) ds = L(t, x)$ in \mathbf{L}^2 . In addition $L(t, x)$ has the representation $L(t, x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_0^t e^{iu(\Lambda_{\alpha,H}(s)-x)} ds du$.*

Next we recall the result on the weak convergence of the partial sum process $S_{[nt]}$. Recall that $\xi_j, -\infty < j < \infty$, is a sequence of i.i.d. random variables belonging to the domain of attraction of a strictly stable law with index $0 < \alpha \leq 2$. For the later purpose we mention that this in particular means

$$E [e^{iu\xi_1}] = \begin{cases} e^{-|u|^\alpha G(|u|)(1+i\beta \text{sign}(u) \tan(\frac{\pi\alpha}{2}))} & \text{if } \alpha \neq 1 \\ e^{-|u|G(|u|)} & \text{if } \alpha = 1 \end{cases} \quad (3)$$

with $|\beta| \leq 1$, where $G(u)$ is slowly varying as $u \rightarrow 0$. In addition, if one lets

$$b_n^{-1} = \inf \{u > 0 : u^\alpha G(u) = n^{-1}\}, \quad (4)$$

then b_n is of the form $n^{1/\alpha} h(n)$ for some slowly varying $h(n)$; in fact $b_n^\alpha \sim nG(b_n^{-1})$. (For the details of these facts, see for instance Bingham et al (1987, page 344.))

Now recall that $c_j, j = 0, 1, \dots$, is a sequence of real numbers such that $c_0 = 1$, and $X_k = \sum_{j=0}^{\infty} c_j \xi_{k-j}$. We impose the following (mutually exclusive) conditions.

(C.1):

$$c_j = j^{H-1-1/\alpha} R(j), \quad \text{with } H \neq 1/\alpha, 0 < H < 1, \quad (5)$$

where $R(j)$ is slowly varying at infinity, and

$$\sum_{j=0}^{\infty} c_j = 0, \quad \text{when } H - 1/\alpha < 0.$$

(C.2):

$$\sum_{j=0}^{\infty} |c_j|^\tau < \infty \quad \text{for some } \tau \text{ such that } 0 < \tau < \alpha, \quad \tau \leq 1,$$

and

$$\sum_{j=0}^{\infty} c_j \neq 0.$$

Throughout what follows we let, with b_n as in (4),

$$\gamma_n = \begin{cases} n^{H-1/\alpha} R(n) b_n & \text{if the condition (C.1) is satisfied} \\ (\sum_{j=0}^{\infty} c_j) b_n & \text{if the condition (C.2) is satisfied.} \end{cases}$$

Because $b_n = n^{1/\alpha} h(n)$, one has $n^{H-1/\alpha} R(n) b_n = n^H u(n)$ for a slowly varying $u(n)$.

The following result is taken from Kasahara and Maejima (1988, Theorems 5.1, 5.2 and 5.3), but see also the references given there, especially Astrauskas (1983).

Proposition 1. (i): *Assume that the condition (C.1) is satisfied. Then the finite dimensional distributions of the process $\gamma_n^{-1} S_{[nt]}$ converge in distribution to those of the LFSM process $\Lambda_{\alpha,H}(t)$, $H \neq 1/\alpha$.*

(ii): *Assume that the condition (C.2) is satisfied. Then the finite dimensional distributions of the process $\gamma_n^{-1} S_{[nt]}$ converge in distribution to those of the α -stable Levy motion $Z_\alpha(t)$*

Note that the Statement (ii) of this result holds for the entire range $0 < \alpha \leq 2$ but our interest is only on the range $1 < \alpha \leq 2$ because as was mentioned earlier the local time for $Z_\alpha(t)$ does not exist when $0 < \alpha \leq 1$. So in the situation of (5) with $H - 1/\alpha < 0$ but $\sum_{j=0}^{\infty} c_j \neq 0$ (which case was excluded in the Statement (i)) this restricts α to either $1 < \alpha < \min\{2, \frac{1}{H}\}$ or $\alpha = 2$ when $\frac{1}{H} > 2$.

3 Statements and discussions of the main results

It is assumed that the constants $\beta_n, n \geq 1$, involved throughout below is such that $\beta_n \rightarrow \infty$ with $n^{-1}\beta_n \rightarrow 0$ as $n \rightarrow \infty$. For any function $h(y)$ we define

$$M_{h,\eta}(y) = \sup\{h(u) : |u - y| \leq \eta\} \quad \text{and} \quad m_{h,\eta}(y) = \inf\{h(u) : |u - y| \leq \eta\}. \quad (6)$$

Also, under the condition (C.2), we shall henceforth restrict to the situation $1 < \alpha \leq 2$ for the reason mentioned above, so that according to our convention the limit $Z_\alpha(t)$ in the Statement (ii) of Proposition 1 becomes $\Lambda_{\alpha,H}(t)$ with $H = 1/\alpha$.

Theorem 2. *Assume that either one of the conditions (C.1) or (C.2) is satisfied. In addition assume that ξ_1 satisfies the Cramér's condition $\limsup_{|u| \rightarrow \infty} |E[e^{iu\xi_1}]| < 1$. Let*

$f(y)$ be such that $M_{|f|,\eta}(y)$ and $M_{f^2,\eta}(y)$ are Lebesgue integrable for some $\eta > 0$ and

$$\int_{-\infty}^{\infty} (M_{f,\delta}(y) - m_{f,\delta}(y)) dy \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad (7)$$

Then the finite dimensional distributions of the process $n^{-1}\beta_n \sum_{k=1}^{[nt]} f(\beta_n(\gamma_n^{-1}S_k + x))$ (indexed by (t, x)) converge in distribution to those of $\left(\int_{-\infty}^{\infty} f(y)dy\right) L(t, -x)$.

Note that $L(t, -x)$ will have the same distribution as that of $L(t, x)$ only when the α -stable Levy motion $Z_\alpha(t)$ involved in the definition of LFSM is symmetric around zero (which is always true in the case of FBM, the case $\alpha = 2$).

Remark 1. There are alternative requirements on $f(y)$ that will imply those stated in Theorem 2. For example one possibility is to assume that the set of discontinuity points of $f(y)$ is of Lebesgue measure zero together with the Lebesgue integrability of $M_{|f|,\eta}(y)$. (It is clear that the condition (7) is then implied by the dominated convergence theorem). Also, as will be indicated later (see the Remark 7 in Section 6 below), it is possible to relax the Lebesgue integrability of $M_{f^2,\eta}(y)$ to that of local Lebesgue integrability. Thus the second possibility is to assume the local Riemann integrability of $f(y)$ together with the Lebesgue integrability of $M_{|f|,\eta}(y)$. In particular Theorem 2 holds for the important situation in which $f(y) = I_{(c,d)}(y)$. (Here the limit will remain the same if the open interval (c, d) is replaced by the closed interval $[c, d]$ or by a semi open interval.)

Remark 2. In the i.i.d. situation $S_k = \sum_{j=1}^k \xi_j$, with $\beta_n = \gamma_n = b_n = \sqrt{n}$, and when $f(y)$ is assumed to be Riemann integrable such that $|f(y)| \leq B(1 + |y|^{-1-c})$ for some $B > 0$ and $c > 0$, Borodin and Ibragimov (1995, Theorem 2.1, Chapter IV, page 143, and Theorem 2.2, Chapter IV, page 145) show that the conclusion of Theorem 2 holds without the Cramér's condition. When $f(y)$ is as above, we mention, without going into the details, the following partial extensions. (a): Theorem 2 extends to the situation where ξ_1 is nonlattice without satisfying the Cramér's condition but only under the restriction $b_n^{-1}\beta_n = O(1)$, where b_n is as in (4). (In the important case $\beta_n = \gamma_n$ this will include the situation of condition (C.2), as well as that of condition (C.1) when $H - 1/\alpha < 0$ but unfortunately will exclude the case $H - 1/\alpha > 0$.) (b): Similarly, Theorem 2 extends to the situation where ξ_1 has a lattice distribution when $b_n^{-1}\beta_n \rightarrow 0$ (which in the case $\beta_n = \gamma_n$ will include the situation of condition (C.1) when $H - 1/\alpha < 0$ but not otherwise).

Remark 3. Regarding the results available in the direction of this paper, Akonom (1993) deals with the situation (of a Brownian motion limit) where $\sum_{j=0}^{\infty} j|c_j| < \infty$ with $\sum_{j=0}^{\infty} c_j \neq 0$, $E[|\xi_1|^p] < \infty$ for some $p > 2$, $\lim_{u \rightarrow \infty} u^\tau E[e^{iu\xi_1}] = 0$ for some $\tau > 0$ and the distribution of ξ_1 has a Lebesgue density. Then the main result obtained there implies the

conclusion of Theorem 2 when $f(y) = I_{[a,b]}(y)$ and $\gamma_n = \sqrt{n}$. This situation is a special case of that of condition (C.2) for $\alpha = 2$, and Theorem 2 requires only that ξ_1 satisfies the Cramér's condition and is in the domain of attraction of a normal distribution and that $\sum_{j=0}^{\infty} |c_j| < \infty$ with $\sum_{j=0}^{\infty} c_j \neq 0$. The underlying situation (with a Brownian motion limit) of Park and Phillips (1999, 2001) is the same as that of Akonom (1993) but the form of $f(y)$ is not restricted to $f(y) = I_{[a,b]}(y)$, though however the (Lipschitz type) conditions imposed there on $f(y)$ are unfortunately rather strong in addition to requiring further moment conditions on ξ_1 .

Tyurin and Phillips (1999) considers the situation where X_k is in the truncated form $X_k = \sum_{j=0}^k c_j \xi_{k-j}$ in addition to the restriction $H - 1/2 < 0$. In addition to the underlying assumptions of Akonom (1993) on the distribution of ξ_1 indicated above, it is further required that $E[|\xi_1|^q] < \infty$ for $q > \frac{1}{H}$. The limiting Gaussian process involved here will be different from a FBM, but will have similar properties. (It is easy to see that the results of the present paper hold for the truncated case $X_k = \sum_{j=0}^k c_j \xi_{k-j}$ also with the changes in the limiting forms taken into account.) \square

Now, an example given in Borodin and Ibragimov (1995, Chapter IV, page 143) shows that the requirement (7) on $f(y)$ in Theorem 2 cannot be avoided entirely. The next result relaxes that requirement but assumes conditions stronger than the Cramér's condition.

Theorem 3. *Assume that either one of the conditions (C.1) or (C.2) is satisfied.*

(i): *Suppose that for some integer n_0 the n_0 -fold convolution of the distribution of ξ_1 has a non-zero absolutely continuous component. Let $f(y)$ be Lebesgue integrable such that $\sup_{y \in \mathbb{R}} |f(y)| < \infty$. Then the conclusion of Theorem 2 holds.*

(ii): *Suppose that, for some integer $n_0 > 0$, $\int |E[e^{iu\xi_1}]|^{n_0} du < \infty$ and $n^{-1}\beta_n \sum_{k=1}^{n_0-1} f(\beta_n(\gamma_n^{-1}S_k + x))$ converges in probability to 0. Assume further that $f(y)$ and $f^2(y)$ are Lebesgue integrable. Then the conclusion of Theorem 2 holds.*

Note that the requirements on $f(y)$ in the first statement is stronger than those in the second statement; consider for instance $f(y) \sim |y|^\tau$ as $|y| \rightarrow 0$ with $0 > \tau > -1/2$. Also, as in Theorem 2 (see Remark 1), the integrability of $f^2(y)$ in the second statement can be relaxed to that of local integrability.

Let us make some remarks regarding the requirement on $n^{-1}\beta_n \sum_{k=1}^{n_0-1} f(\beta_n(\gamma_n^{-1}S_k + x))$ in the Statement (ii). First, it is redundant when $n_0 = 1$. When $x = 0$ and $\beta_n = \gamma_n$, an important case in applications, the requirement is satisfied because the quantity reduces to $n^{-1}\beta_n \sum_{k=1}^{n_0-1} f(S_k)$ which clearly converges in probability to 0 in view of $n^{-1}\beta_n \rightarrow 0$.

Suppose that $x \neq 0$. Then the additional condition $\lim_{a \rightarrow \infty} \sup_{|y| \geq a} |f(y)| < \infty$ is

sufficient because, for each $1 \leq k < n_0$, with probability tending to one $\beta_n (\gamma_n^{-1} S_k + x)$ will be supported in a neighborhood of $\pm\infty$. The same is the case when $x = 0$ and $\beta_n \gamma_n^{-1} \rightarrow \infty$.

In the remaining case $x = 0$ and $\beta_n \gamma_n^{-1} \rightarrow 0$, suppose for instance that $f(y) \sim |y|^\tau$ as $|y| \rightarrow 0$ with $0 > \tau > -1/2$. Then with probability tending to one, $n^{-1} \beta_n |f(\beta_n \gamma_n^{-1} S_k)| \leq C n^{-1} \beta_n |\beta_n \gamma_n^{-1}|^\tau = C (n^{-1} \beta_n)^{1+\tau} (n \gamma_n^{-1})^\tau \rightarrow 0$.

Remark 4. So far our results are for $f_n(y) = \beta_n f(\beta_n y)$ based on $f(y)$. It is possible to extend Theorems 2 and 3 to more general $f_n(y)$ that satisfy the conditions: $\sup_n \int |f_n(y)| dy < \infty$, $\sup_n \frac{1}{n} \int_{-\infty}^{\infty} |f_n^2(y)| dy \rightarrow 0$, $\lim_{\kappa \rightarrow \infty} \sup_n \int_{\{|y| \geq \kappa\}} |f_n(y)| dy = 0$ and, letting

$$F_n(y) = \begin{cases} \int_0^y f_n(u) du & \text{if } y \geq 0 \\ -\int_y^0 f_n(u) du & \text{if } y < 0, \end{cases}$$

there is an $F(y)$ such that $F_n(y) \rightarrow F(y)$ at all *continuity points* of $F(y)$. These conditions are satisfied when $f_n(y) = \beta_n f(\beta_n y)$ with $F(y) = \int_0^\infty f(u) du$ if $y \geq 0$ and $F(y) = \int_{-\infty}^0 f(u) du$ if $y < 0$, and in fact we shall use the assumptions on $f(y)$ only in the form of the above conditions. Part two of Theorem 3 extends as follows. Assume, for some integer $n_0 > 0$, $\int |E[e^{iu\xi_1}]|^{n_0} du < \infty$ and $n^{-1} \sum_{k=1}^{n_0-1} f_n(\gamma_n^{-1} S_k + x)$ converges in probability to 0. Then, if $f_n(y)$ satisfy the above conditions, $n^{-1} \sum_{k=1}^{[nt]} f_n(\gamma_n^{-1} S_k + x)$ converges in distribution to $\int_{-\infty}^{\infty} L(t, y-x) dF(y)$. Similarly part one of Theorem 3 extends, under the additional assumption $\sup_{\{n,y\}} \sigma_n^{-1} |f_n(y)| < \infty$ for some $\sigma_n \rightarrow \infty$ with $\frac{\sigma_n}{n} \rightarrow 0$.

The next result may be viewed as a discrete approximation to the local time of the LFSM, which we obtain as a by-product to the Statement (ii) of Theorem 3 (whose requirements are satisfied with $n_0 = 1$). Note that the approximation is in \mathbf{L}^2 , in contrast to the distributional approximation in Theorems 2 and 3. Note also that in this case $\gamma_n = n^H$.

Theorem 4. *Suppose that $f(y)$ and $f^2(y)$ are Lebesgue integrable. Then, for each t ,*

$$\frac{\beta_n}{n} \sum_{k=1}^{[nt]} f(\beta_n (\Lambda_{\alpha,H}(k/n) - x)) \rightarrow \left(\int_{-\infty}^{\infty} f(y) dy \right) L(t, x) \quad \text{in } \mathbf{L}^2.$$

The next result is a continuous analogue of the preceding result.

Theorem 5. *Assume that $f(y)$ and $f^2(y)$ are Lebesgue integrable. Then*

$$\frac{1}{\kappa^{1-H}} \int_0^{\kappa t} f(\kappa^H (\Lambda_{\alpha,H}(s/\kappa) - x)) ds \rightarrow \left(\int_{-\infty}^{\infty} f(y) dy \right) L(t, x) \quad \text{in } \mathbf{L}^2$$

for each t and x as $\kappa \rightarrow \infty$.

As noted in connection with the Statement (ii) of Theorem 3, the Lebesgue integrability of $f^2(y)$ in Theorems 4 and 5 can be relaxed to that of local Lebesgue integrability.

Remark 5. Note that because the distribution of $n^H \Lambda_{\alpha,H}(k/n)$ is the same as that of $\Lambda_{\alpha,H}(k)$, it follows from Theorem 4 that $\frac{1}{n^{1-H}} \sum_{k=1}^{[nt]} f(\Lambda_{\alpha,H}(k) - xn^H)$ converges in distribution to $\left(\int_{-\infty}^{\infty} f(y)dy\right) L(t, x)$ as $n \rightarrow \infty$. Similarly $\frac{1}{\kappa^{1-H}} \int_0^{\kappa t} f(\Lambda_{\alpha,H}(s) - x\kappa^H) ds$ converges in distribution to $\left(\int_{-\infty}^{\infty} f(y)dy\right) L(t, x)$ as $\kappa \rightarrow \infty$, in view of Theorem 5.

Remark 6. One has $\frac{1}{\kappa^{1-H}} \int_0^{\kappa t} f(\kappa^H(\Lambda_{\alpha,H}(s/\kappa) - x)) ds = \kappa^H \int_0^{\kappa t} f(\kappa^H(\Lambda_{\alpha,H}(s) - x)) ds$. When $f(y) = \mathbb{I}_{[0,1)}(y)$ and $\kappa^H = 1/\eta$ the right hand side reduces to $\frac{1}{\eta} \int_0^t \mathbb{I}_{[x, x+\eta)}(\Lambda_{\alpha,H}(s)) ds$.

4 Proofs of the results

The Proposition 6 stated next is just a formalization of the approximation (1), the proof of which is postponed to the next two sections because it requires many auxiliary results.

In this section we derive Theorems 0 and 2-5 as consequences of it. Recall that we let $\phi_\varepsilon(z) = \frac{1}{\varepsilon\sqrt{2\pi}} \exp\left\{-\frac{z^2}{2\varepsilon^2}\right\}$ and $\phi(z) = \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{z^2}{2}\right\}$.

Proposition 6. Let $f_n(y) = \beta_n f(\beta_n y)$. Assume that the conditions of any one of Theorems 2 or 3 are satisfied. Let $L_n(t, x) = \frac{1}{n} \sum_{k=q}^{[nt]} f_n(\gamma_n^{-1} S_k + x)$, where $q = 1$ except under the second part of Theorem 3 in which case $q = n_0$ with n_0 as involved there, and

$$L_{n,\varepsilon}(t, x) = \frac{1}{n} \sum_{k=1}^{[nt]} \int_{-\infty}^{\infty} f_n(\gamma_n^{-1} S_k + x + z\varepsilon) \phi(z) dz.$$

Then

$$\lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \lim_{t,x} E[(L_n(t, x) - L_{n,\varepsilon}(t, x))^2] = 0.$$

We next deal with $L_{n,\varepsilon}(t, x)$ through the following series of steps.

Lemma 7. For each $\varepsilon > 0$,

$$\sup_{t,x} \left| L_{n,\varepsilon}(t, x) - \left(\int_{-\infty}^{\infty} f(z) dz \right) \frac{1}{n} \sum_{k=1}^{[nt]} \phi_\varepsilon(\gamma_n^{-1} S_k + x) \right| \quad (8)$$

is bounded by a nonrandom quantity that converges to 0 as $n \rightarrow \infty$.

Proof. For convenience take $\varepsilon = 1$. Let $F_n(y) = \int_{-\infty}^y \beta_n f(\beta_n u) du$, and define $F(y)$ such that $F(y) = \int_{-\infty}^{\infty} f(u) du$ if $y \geq 0$ and $F(y) = 0$ if $y < 0$. Note that $F_n(y) \rightarrow F(y)$ at all continuity points of $F(y)$ and $F(b) - F(a) = 0$ if $0 \notin (a, b]$. Now $L_{n,\varepsilon}(t, x)$ takes the form $\int_{-\infty}^{\infty} \left(\frac{1}{n} \sum_{k=1}^{[nt]} \phi(y - \gamma_n^{-1} S_k - x) \right) dF_n(y)$. The difference between this and

$$\int_{\{|y| \leq \nu\}} \left(\frac{1}{n} \sum_{k=1}^{[nt]} \phi(y - \gamma_n^{-1} S_k - x) \right) dF_n(y) \quad (9)$$

is bounded in absolute value by $C \int_{\{|y|>\nu\}} d|F_n|(y) = C \int_{\{|u|>\beta_n\nu\}} |f(u)| du$. Define $y_{m,i}$, $i = -m, \dots, 0, \dots, m$ such that $y_{m,-m} = -\nu < y_{m,-m+1} < \dots < y_{m,m-1} < y_{m,m} = \nu$ and $\sup_i |y_{m,i} - y_{m,i-1}| \leq \frac{2\nu}{m}$. Then the difference between (9) and

$$\sum_{i=-m}^{m-1} \left(\frac{1}{n} \sum_{k=1}^{[nt]} \phi(y_{m,i} - \gamma_n^{-1} S_k - x) \right) \int_{y_{m,i}}^{y_{m,i+1}} dF_n(y) \quad (10)$$

is bounded in absolute value by $C \frac{\nu}{m} \int_{\{|y| \leq \nu\}} d|F_n|(y) \leq C \frac{\nu}{m}$. Further the difference between (10) and $\sum_{i=-m}^{m-1} \left(\frac{1}{n} \sum_{k=1}^{[nt]} \phi(y_{m,i} - \gamma_n^{-1} S_k - x) \right) \int_{y_{m,i}}^{y_{m,i+1}} dF(y)$ is bounded in absolute value by $C \sum_{i=-m}^{m-1} \left| \int_{y_{m,i}}^{y_{m,i+1}} d(F_n(y) - F(y)) \right|$. Thus it follows that (8) is bounded in absolute value by a constant multiple of $\int_{\{|u|>\beta_n\nu\}} |f(u)| du + \frac{\nu}{m} + \sum_{i=-m}^{m-1} \left| \int_{y_{m,i}}^{y_{m,i+1}} d(F_n(y) - F(y)) \right|$. Denote this by $Q(\nu, m, n)$. It is clear that $\lim_{\nu \rightarrow \infty} \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} Q(\nu, m, n) = 0$. \square

Lemma 8. *For each $\varepsilon > 0$, the finite dimensional distributions of $\frac{1}{n} \sum_{k=1}^{[nt]} \phi_\varepsilon(\gamma_n^{-1} S_k + x)$ (indexed by (t, x)) converge in distribution to those of $\int_0^t \phi_\varepsilon(\Lambda_{\alpha, H}(s) + x) ds$.*

Proof. For notational convenience take $\varepsilon = 1$. Also we take $x = 0$ for simplicity so that we consider the process $n^{-1} \sum_{k=1}^{[nt]} \phi(\gamma_n^{-1} S_k)$ indexed only by t . We now invoke Gikhman and Skorokhod (1969, Theorem 1, page 485). According to this result, letting $\zeta_n(s) = \phi(\gamma_n^{-1} S_{[ns]})$, we need to check that (the finite dimensional distributions of $\zeta_n(s)$ being convergent to those of the limit $\zeta(s) = \phi(\Lambda_{\alpha, H}(s))$ by Proposition 1)

$$\sup_{s, n} E[|\zeta_n(s)|] < \infty, \quad \lim_{\eta \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_{|s_1 - s_2| \leq \eta} E[|\zeta_n(s_1) - \zeta_n(s_2)|] = 0.$$

Because $\lim_{\kappa \rightarrow \infty} \sup_{|y| > \kappa} \phi(y) = 0$, it is enough to verify that, for every $\delta > 0$,

$$\lim_{\eta \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_{|s_1 - s_2| \leq \eta} P(|\gamma_n^{-1} S_{[ns_1]} - \gamma_n^{-1} S_{[ns_2]}| > \delta) = 0$$

Note that $\sup_{|s_1 - s_2| \leq \eta} P(|\gamma_n^{-1} S_{[ns_1]} - \gamma_n^{-1} S_{[ns_2]}| > \delta) = \sup_{0 < \tau \leq \eta} P(|\gamma_n^{-1} S_{[n\tau]}| > \delta)$ by the stationarity of $\{X_k, k \geq 1\}$. (Here the stationarity is used only for convenience and can be avoided.) Now, Theorems 2.2 and 4.1 in Kasahara and Maejima (1988) give bounds for $\sup_{0 < \tau \leq \eta} P(|\gamma_n^{-1} S_{[n\tau]}| > \delta)$, respectively when $0 < \alpha < 2$ and when $\alpha = 2$. Using the arguments similar to those used in Kasahara and Maejima (1988, Section 5), these bounds converge to zero by first letting $n \rightarrow \infty$ and then $\eta \rightarrow 0$. This completes the proof. \square

The next result is related to Theorem 4.

Lemma 9. *For each $\varepsilon > 0$ and for all t and x , $n^{-1} \sum_{k=1}^{[nt]} \phi_\varepsilon(\Lambda_{\alpha, H}(\frac{k}{n}) + x) \rightarrow \int_0^t \phi_\varepsilon(\Lambda_{\alpha, H}(s) + x) ds$ in \mathbf{L}^2 .*

Proof. t, x and ε being fixed, take for notational simplicity $t = 1, x = 0$ and $\varepsilon = 1$. Write $n^{-1} \sum_{k=1}^n \phi(\Lambda_{\alpha, H}(\frac{k}{n}))$ in the form $\int_0^1 \phi\left(\Lambda_{\alpha, H}\left(\frac{[ns]}{n}\right)\right) ds$. Then the proof is clear

because $\Lambda_{\alpha,H} \left(\frac{[ns]}{n} \right) - \Lambda_{\alpha,H}(s)$ converges to 0 in probability (see Samorodnitsky and Taqqu (1994, Proposition 7.4.3)) and because $\sup_z \phi(z) < \infty$. \square

The next result is a continuous analogue of Proposition 6 for the LFSM situation.

Proposition 10. *Suppose that $f(y)$ and $f^2(y)$ are Lebesgue integrable. Let $L_\kappa^*(t, x) = \frac{1}{\kappa^{1-H}} \int_0^{\kappa t} f(\kappa^H (\Lambda_{\alpha,H}(s/\kappa) - x)) ds$. Further let $L_\varepsilon(t, x) = \left(\int_{-\infty}^{\infty} f(y) dy \right) \int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds$. Then*

$$\lim_{\varepsilon \rightarrow 0} \limsup_{\kappa \rightarrow \infty} \sup_{x,t} E [|L_\kappa^*(t, x) - L_\varepsilon(t, x)|^2] \rightarrow 0.$$

The proof of this result is essentially contained in the proof of Proposition 6 for the situation of the second part (with $n_0 = 1$) of Theorem 3. The next proposition will be the only remaining fact required (apart from the proof of Proposition 6) to complete the proofs of Theorems 0 and 2 - 5.

Proposition 11. *For each t and x , $\int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds \rightarrow L(t, x)$ in \mathbf{L}^2 as $\varepsilon \downarrow 0$, where $L(t, x)$ is the local time of $\Lambda_{\alpha,H}(s)$, that is, satisfies the requirement (2). In addition $L(t, x)$ has the representation of Theorem 0, that is $L(t, x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_0^t e^{iu(\Lambda_{\alpha,H}(s) - x)} ds du$.*

Proof. Let $\Upsilon_\varepsilon = \int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds$. We first show that Υ_ε is a Cauchy sequence in \mathbf{L}^2 . For notational convenience, we restrict for the moment to $t = 1$ and suppress the occurrence of x . We need to show that $E [(\Upsilon_{\varepsilon_1} - \Upsilon_{\varepsilon_2})^2] \rightarrow 0$ as $(\varepsilon_1, \varepsilon_2) \downarrow 0$. Note that

$$e^{-v^2/2\varepsilon^2} = \frac{\varepsilon}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ivu} e^{-u^2\varepsilon^2/2} du = \frac{\varepsilon}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ivu} e^{-u^2\varepsilon^2/2} du.$$

Hence, because $E[\Upsilon_{\varepsilon_1} \Upsilon_{\varepsilon_2}] = E \left[\int_0^1 \int_0^1 \phi_{\varepsilon_1}(\Lambda_{\alpha,H}(s_1)) \phi_{\varepsilon_2}(\Lambda_{\alpha,H}(s_2)) ds_1 ds_2 \right]$,

$$E[\Upsilon_{\varepsilon_1} \Upsilon_{\varepsilon_2}] = \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E \left[\int_0^1 \int_0^1 e^{iu\Lambda_{\alpha,H}(s_1) - iv\Lambda_{\alpha,H}(s_2)} ds_1 ds_2 \right] e^{-\frac{u^2\varepsilon_1^2}{2} - \frac{v^2\varepsilon_2^2}{2}} dudv. \quad (11)$$

Here the order of integration is interchanged, which is permissible for each fixed ε_1 and ε_2 , as can be seen using the fact $\int_0^1 \int_0^1 |E[e^{iu\Lambda_{\alpha,H}(s_1) - iv\Lambda_{\alpha,H}(s_2)}]| ds_1 ds_2 \leq 1$. Now if

$$\frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E \left[\int_0^1 \int_0^1 e^{iu\Lambda_{\alpha,H}(s_1) - iv\Lambda_{\alpha,H}(s_2)} ds_1 ds_2 \right] dudv < \infty, \quad (12)$$

then it is clear from (11) that $E[\Upsilon_{\varepsilon_1} \Upsilon_{\varepsilon_2}]$ converges to (12) as $(\varepsilon_1, \varepsilon_2) \rightarrow 0$ by the dominated convergence theorem. (Note that $E \left[\int_0^1 \int_0^1 e^{iu\Lambda_{\alpha,H}(s_1) - iv\Lambda_{\alpha,H}(s_2)} ds_1 ds_2 \right] \geq 0$.) The same is also true for $E[\Upsilon_{\varepsilon_1}^2]$ and $E[\Upsilon_{\varepsilon_2}^2]$, so that $E[(\Upsilon_{\varepsilon_1} - \Upsilon_{\varepsilon_2})^2] \rightarrow 0$ as $(\varepsilon_1, \varepsilon_2) \downarrow 0$.

We now verify (12), in a stronger form that will be needed below. Note that, making now the occurrence of x explicit, for the $L_\kappa^*(t, x)$ defined in Proposition 10, one can directly check that $\sup_\kappa \sup_x E[(L_\kappa^*(t, x))^2] < \infty$ (See Remark 8 of Section 6.) Thus in view of

Proposition 10, it follows that $\limsup_{\varepsilon \rightarrow 0} \sup_x E \left[\left(\int_0^1 \phi_\varepsilon(\Lambda_{\alpha,H}(s)) ds \right)^2 \right] < \infty$. Hence, in view of the identity (11) (with $\varepsilon_1 = \varepsilon_2$), it follows by Fatou's Lemma that

$$\sup_x \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left[\int_0^1 \int_0^1 E \left[e^{iu(\Lambda_{\alpha,H}(s_1)-x)-iv(\Lambda_{\alpha,H}(s_2)-x)} \right] ds_1 ds_2 \right] dudv < \infty. \quad (13)$$

Actually we have shown that (11) converges to (12) because (11) is bounded above by (12).

Thus we have established the Cauchy convergence of $\int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds$ in \mathbf{L}^2 , which entails the convergence in \mathbf{L}^2 . Denote the limit by $L(t, x)$. In particular, (13) gives

$$\int_c^d \int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds dx \rightarrow \int_c^d L(t, x) dx \quad \text{in } \mathbf{L}^1 \quad (14)$$

for every interval $[c, d]$. It remains to show that the limit $L(t, x)$ is indeed a local time, that is, (2) holds. It is sufficient to verify that $\int_0^t \mathbb{I}_{[c,d]}(\Lambda_{\alpha,H}(s)) ds = \int_c^d L(t, x) dx$ for every interval $[c, d]$. In view of Remark 6, Proposition 10 in particular entails that

$$\frac{1}{\eta} \int_0^t \mathbb{I}_{[x, x+\eta]}(\Lambda_{\alpha,H}(s)) ds - \int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds \rightarrow 0 \quad \text{in } \mathbf{L}^2$$

as $\eta \rightarrow 0$ first and then $\varepsilon \rightarrow 0$, uniformly in x (in the sense of Proposition 10). Hence

$$\int_0^t \mathbb{I}_{[c,d]}(\Lambda_{\alpha,H}(s)) ds - \int_c^d \int_0^t \phi_\varepsilon(\Lambda_{\alpha,H}(s) - x) ds dx \rightarrow 0 \quad \text{in } \mathbf{L}^2$$

as $\varepsilon \rightarrow 0$. In view of (14), this completes the proof. \square

Now note that Propositions 10 and 11 together with Remark 6 proves Theorem 0 of Section 2. In the same way Theorems 2 and 3 follow from Propositions 6 and 11 and Lemmas 7 and 8. Thus it only remains to establish Proposition 6.

5 Auxiliary results for the proof of Proposition 6

We first obtain some estimates on the behavior of the characteristic function of

$$S_j^* = \gamma_j^{-1} \sum_{k=1}^j g(k) \xi_{j-k}. \quad (15)$$

(The reason for considering S_j^* , which is in the form of a finite order sum of independent r.v.s, will become clear in the next section.) Here $g(k) = \sum_{i=0}^k c_i$ with c_i as in the conditions (C.1) or (C.2). Only the distributional properties of S_j^* will be required so that for convenience we take $S_j^* = \gamma_j^{-1} \sum_{k=1}^j g(k) \xi_k$. Under the condition (C.1) we have $c_j = j^{H-1-1/\alpha} R(j)$ for a slowly varying $R(j)$ (with $\sum_{i=0}^{\infty} c_i = 0$ if $H - 1/\alpha < 0$). For simplicity we shall restrict to the situation $c_j \sim C j^{H-1-1/\alpha}$, $j \rightarrow \infty$, for a suitable constant

C . Then note that $g(k) = \sum_{i=1}^k c_i \sim C' k^{H-1/\alpha}$, $k \rightarrow \infty$, for some constant C' , so that for convenience S_j^* can be taken to be of the form (with b_j as in (4)).

$$S_j^* = \sum_{k=1}^j \binom{k}{j}^{H-1/\alpha} \frac{\xi_k}{b_j}. \quad (16)$$

Note that in view of (3) there is a $\delta > 0$ such that, letting $\psi(v) = E[e^{iv\xi_1}]$,

$$|\psi(v)| \leq e^{-|v|^\alpha G(|v|)} \quad \text{for all } |v| \leq \delta \quad (17)$$

where $G(w)$ is slowly varying at 0.

In the rest of this section it is assumed without further mentioning that either one of the conditions (C.1) or (C.2) is satisfied. Further the constant b_j is as in (4).

Lemma 12. *Let $\widehat{H}_j(u)$ be the characteristic function of S_j^* defined in (15). Then there are constants $A > 0, \lambda > 0, d > 0$ and $0 < c < \alpha$ such that $|\widehat{H}_j(u)| \leq Ae^{-d|u|^c}$ for all $|u| \leq \lambda b_j$ and $j \geq 1$, where b_j is as in (4).*

Proof. It is enough to prove the result for all sufficiently large j , because for any $j_0 > 1$

$$\left| \widehat{H}_j(u) \right| \leq 1 = e^{d|\lambda b_{j_0}|^c} e^{-d|\lambda b_{j_0}|^c} \leq e^{d|\lambda b_{j_0}|^c} e^{-d|u|^c} \quad \text{for } 1 \leq j \leq j_0 \text{ and } |u| \leq \lambda b_j.$$

In the same way it is enough to consider u such that $|u| \geq C$ for some $C > 0$. We have $|\widehat{H}_j(u)| = \prod_{k=1}^j |\psi(\gamma_j^{-1} g(k)u)|$. Consider the situation of condition (C.1). Then (taking into account the simplification (16)), $|\widehat{H}_j(u)| = \prod_{k=1}^j \left| \psi \left(\left(\frac{k}{j} \right)^{H-1/\alpha} \frac{u}{b_j} \right) \right|$. Suppose first that $H - 1/\alpha > 0$. Then $\left| \left(\frac{k}{j} \right)^{H-1/\alpha} \frac{u}{b_j} \right| \leq \delta$ for all $|u| \leq \delta b_j$. Thus in view of (17)

$$\left| \widehat{H}_j(u) \right| \leq \exp \left\{ - \sum_{k=1}^j \left| \left(\frac{k}{j} \right)^{H-1/\alpha} \frac{u}{b_j} \right|^\alpha G \left(\left(\frac{k}{j} \right)^{H-1/\alpha} \frac{|u|}{b_j} \right) \right\} \quad \text{for all } |u| \leq \delta b_j.$$

Recall that $b_j^\alpha \sim jG(b_j^{-1})$ (the $G(u)$ in (4) and that in (17) being the same). One can assume for convenience that $b_j^\alpha = jG(b_j^{-1})$. Then the sum in the preceding exponent becomes

$$|u|^\alpha \frac{1}{j} \sum_{k=1}^j \left| \left(\frac{k}{j} \right)^{H-1/\alpha} \right|^\alpha \frac{1}{G(b_j^{-1})} G \left(\left(\frac{k}{j} \right)^{H-1/\alpha} \frac{|u|}{b_j} \right). \quad (18)$$

Note that $G(w)$ being continuous it is bounded on compacts. Hence by Potter's inequality (see Bingham et al (1987, Statement (ii) of Theorem 1.5.6, page 25), for every $\delta > 0$ there is a $B > 0$ such that $|G(x)/G(y)| \leq B \max\{(x/y)^\delta, (x/y)^{-\delta}\}$ for all $x > 0, y > 0$. Hence it follows easily that for a given $\epsilon > 0$ there are $C > 0$ and $C_1 > 0$ such that

$\inf_{k \geq [j\epsilon]} \left| \frac{1}{G(b_j^{-1})} G \left(\left(\frac{k}{j} \right)^{H-1/\alpha} \frac{|u|}{b_j} \right) \right| \geq C_1 |u|^{-\delta}$ for all sufficiently large j and for all $|u| \geq C$.

Thus (18) is bounded below, for some $C' > 0$, by $C' |u|^{\alpha-\delta}$ for all sufficiently large j and for all $|u| \geq C$. This proves the result under the condition (C.1) when $H - 1/\alpha > 0$.

When $H - 1/\alpha < 0$, note that whenever $k \geq [j\epsilon]$ for a given ϵ one has, for some $C > 0$, $\left(\frac{k}{j} \right)^{H-1/\alpha} \leq C$ for all sufficiently large j . Hence essentially the same arguments used above apply for this case also.

It remains to consider the case under the condition (C.2). By assumption $\sum_{i=0}^{\infty} c_i \neq 0$ and $\sum_{i=0}^{\infty} |c_i| < \infty$, so that for some positive constants ϑ and κ one has $\vartheta \leq \left| \sum_{i=0}^k c_i \right| \leq \kappa$ for all sufficiently large k . Further in this case $\gamma_j^{-1} = b_j^{-1} (\sum_{i=0}^{\infty} c_i)^{-1}$. Hence the proof of this case is contained in the above arguments. This completes the proof of the lemma. \square

Lemma 13. *Let $\widehat{H}_j(u)$ be the characteristic function of S_j^* defined in (15). Assume that ξ_1 satisfies the Cramér's condition of Theorem 2. Then for any $d > 0$ there is a $B > 0$ and a $0 < \rho < 1$ such that $\sup_{|u| \geq db_j} \left| \widehat{H}_j(u) \right| \leq B \rho^j$ for all $j \geq 1$, where b_j is as in (4).*

Proof. As in Lemma 12, it is enough to prove the statement for all sufficiently large j . Now recall that the Cramér's condition is actually equivalent to $\sup_{|z| \geq a} |\psi(z)| < 1$ for all $a > 0$. First consider the situation under (C.1). When $H - 1/\alpha < 0$ (and taking into account the simplification (16)), we have $\left(\frac{k}{j} \right)^{H-1/\alpha} \geq 1$ for all $1 \leq k \leq j$. Thus

$$\sup_{|u| \geq db_j} \left| \widehat{H}_j(u) \right| = \sup_{|u| \geq db_j} \prod_{k=1}^j \left| \psi \left(\left(\frac{k}{j} \right)^{H-1/\alpha} \frac{u}{b_j} \right) \right| \leq \left(\sup_{|z| \geq d} |\psi(z)| \right)^j.$$

Now consider $H - 1/\alpha > 0$. Given $\epsilon > 0$, there is a $C > 0$ and a j_0 such that $\left(\frac{k}{j} \right)^{H-1/\alpha} \geq C$ for all $k \geq [j\epsilon]$ and $j \geq j_0$. Hence, as before, $\sup_{|u| \geq db_j} \left| \widehat{H}_j(u) \right| \leq (\sup_{|z| \geq dC} |\psi(z)|)^{j-[j\epsilon]}$, $j \geq j_0$. Because for some $\mu > 0$, $j - [j\epsilon] \geq \mu j$ for all sufficiently large j , this proves the result under (C.1). The proof under (C.2) uses the same arguments. \square

To proceed further we need the following result, contained in Bhattacharya and Ranga Rao (1976, the proof of Lemma 11.1, page 93).

Lemma 14. *Let μ and ν be finite measures on R^k . Let η be a positive number and K_η a probability measure on R^k satisfying $K_\eta(\{x : |x| \leq \eta\}) = 1$. Let h be a real valued Borel measurable function on R^k such that $|M_{h,\eta}(x)|$ and $|m_{h,\eta}(x)|$ are integrable with respect to μ and ν (where $M_{h,\eta}(x)$ and $m_{h,\eta}(x)$ are as defined in (6)). Then*

$$\begin{aligned} \int f d(\mu - \nu) &\geq \int m_{f,\eta} d(\mu - \nu) * K_\eta - \int (f - m_{f,2\eta}) d\nu \\ &\leq \int M_{f,\eta} d(\mu - \nu) * K_\eta + \int (M_{f,2\eta} - f) d\nu. \end{aligned}$$

The K_η above will be chosen such that its characteristic function $\widehat{K}_\eta(u)$ satisfies

$$|\widehat{K}_\eta(u)| \leq C \exp\{-(\eta|u|)^{1/2}\} \quad (19)$$

for all real u , where C is a constant (independent of η). This is possible in view of Bhattacharya and Ranga Rao (1976, Corollary 10.4, page 88), where K_η is used extensively as a smoothing device. Next we give some inequalities that will be used in a rather crucial manner in the next section. For this reason we need to state the bounds explicitly.

Lemma 15. *Assume that ξ_1 satisfies the Cramér's condition. Let K_η be the smoothing probability measure of Lemma 14 satisfying (18). Then for some $0 < \rho < 1$,*

$$\int \left| \widehat{H}_j(u) \right| \left| \widehat{K}_\eta(u) \right| du \leq C (1 + \eta^{-1} \rho^j)$$

and

$$\int \left| e^{-u^2 \varepsilon^2 / 2} - 1 \right| \left| \widehat{H}_j(u) \right| \left| \widehat{K}_\eta(u) \right| du \leq C (\varepsilon^2 + \eta^{-1} \rho^j).$$

Proof. Let $\lambda > 0, \alpha > c > 0, d > 0$ be positive constants involved in Lemma 12 and let $0 < \rho < 1$ be as in Lemma 13. Then, recalling the inequality (19) for $\left| \widehat{K}_\eta(u) \right|$,

$$\begin{aligned} \int \left| \widehat{H}_j(u) \right| \left| \widehat{K}_\eta(u) \right| du &\leq \int_{\{|u| \leq \lambda b_j\}} \left| \widehat{H}_j(u) \right| du + \int_{\{|u| > \lambda b_j\}} \left| \widehat{H}_j(u) \right| \left| \widehat{K}_\eta(u) \right| du \\ &\leq C \int_{\{|u| \leq \lambda b_j\}} e^{-d|u|^c} du + C \rho^j \int_{\{|u| > \lambda b_j\}} e^{-(\eta|u|)^{1/2}} du \end{aligned}$$

which proves the first part. The second part also follows in the same way using the inequality $\left| e^{-u^2 \varepsilon^2 / 2} - 1 \right| \leq u^2 \varepsilon^2 / 2$ over the range $\{|u| \leq \lambda b_j\}$ and using the inequality $\left| e^{-u^2 \varepsilon^2 / 2} - 1 \right| \leq 2$ over the range $\{|u| > \lambda b_j\}$. \square

Lemma 16. *Suppose that for some integer n_0 the n_0 -fold convolution of the distribution of ξ_1 has a non-zero absolutely continuous component. Let H_j be the probability distribution S_j^* defined in (15). Then there are measures H_j^* and H_j^{**} such that $H_j = H_j^* + H_j^{**}$, satisfying the following properties:*

(i) *For every $r > 0$ there is a constant L_r such that $H_j^{**}(R) \leq L_r j^{-r}$ for all $j \geq n_0$.*

(ii) *Let \widehat{H}_j^* be the Fourier transform of H_j^* . Then there are constants $A > 0, \lambda > 0, d > 0$ and $0 < c < \alpha$ such that for every $j \geq n_0$ and for every $r > 0$, $\left| \widehat{H}_j^* \right| \leq A e^{-d|u|^c} + L_r j^{-r}$ for all $|u| \leq \lambda b_j$ and for a suitable constant L_r .*

(iii) *There are constants $B > 0$ and $0 < \rho < 1$ and an integrable function $g(v) \geq 0$ such that $\sup_{|u| \geq \lambda b_j} \left| \widehat{H}_j^*(u) \right| \leq B g(b_j^{-1} u) \rho^j$ for all $j \geq 2n_0$.*

Proof. The proof is similar to the i.i.d. case $j^{-1/2} \sum_{i=1}^j \xi_i$ (see Prohorov (1952) and Le Cam (1960, pages 68-72)). We briefly recall the arguments. For simplicity

we consider the case $n_0 = 1$. First, letting F for the distribution of ξ_1 , one writes $F = \theta G_1 + (1 - \theta) G_2$ where $0 < \theta < 1$ and G_1 and G_2 are probability measures such that G_1 is absolutely continuous with respect to Lebesgue measure with a Lebesgue density uniformly bounded by a constant. (In particular the density of G_1 is square integrable and hence $|\widehat{G}_1|^2$ is also integrable.) This is possible because F is not entirely disjoint from the Lebesgue measure. Then the characteristic function $\psi_F(u)$ of F has the corresponding decomposition $\psi_F(u) = \theta\psi_{G_1}(u) + (1 - \theta)\psi_{G_2}(u)$. Hence $\widehat{H}_j(u) = \prod_{k=1}^j (\theta\psi_{G_1}(\gamma_j^{-1}g(k)u) + (1 - \theta)\psi_{G_2}(\gamma_j^{-1}g(k)u))$ which can be written in the form $\widehat{H}_j(u) = \sum_{l=0}^j \sum_{*l} \theta^l (1 - \theta)^{j-l} \prod_{k=1}^j \psi_{*l}(\gamma_j^{-1}g(k)u)$ where ψ_{*l} is either ψ_{G_1} or ψ_{G_2} such that ψ_{G_1} occurs in the product $\prod_{k=1}^j \psi_{*l}(\gamma_j^{-1}g(k)u)$ exactly l times, and the sum \sum_{*l} is with respect to all such products for a given l . Now let $I_j^c = \{l : |l - j\theta| > j^{3/4}\}$. Then H_j^{**} is the measure corresponding to the Fourier transform $\sum_{l \in I_j^c} \sum_{*l} \theta^l (1 - \theta)^{j-l} \prod_{k=1}^j \psi_{*l}(\gamma_j^{-1}g(k)u)$. It is clear that $H_j^{**}(R)$ is bounded in absolute value by $\sum_{l \in I_j^c} \binom{j}{l} \theta^l (1 - \theta)^{j-l}$, which is known to have the bound stated in the Statement (i). For the Statement (ii) use the fact that $|\widehat{H}_j^*(u)| \leq |\widehat{H}_j(u)| + H_j^{**}(R)$ and then use the bound in Statement (i) for $H_j^{**}(R)$ together with the bound in Lemma 12 for $|\widehat{H}_j(u)|$.

Proof of the Statement (iii) uses essentially the same arguments of the i.i.d. case given in Prohorov (1952) or LeCam (1960) mentioned above, the essential facts being the Cramér's condition for G_1 and the function $g(v)$ in the statement (iii) taken to be $|\widehat{G}_1(av)|^2$ for a suitable constant a , which is integrable as indicated earlier. The necessary modifications needed for the present case being essentially the same as those used in the proofs of the Lemmas 12 and 13 above, the proof is concluded. \square

Lemma 17. *Under the situation of the preceding Lemma 16, for some $0 < \rho < 1$,*

$$\int |\widehat{H}_j^*(u)| du \leq C (1 + \rho^j)$$

and

$$\int |e^{-u^2\varepsilon^2/2} - 1| |\widehat{H}_j^*(u)| du \leq C (\varepsilon^2 + \rho^j)$$

for all $j \geq 2n_0$.

Proof. Regarding the first statement, according to the Statements (ii) and (iii) of Lemma 16, there is a $0 < \rho < 1$ such that for any $r > 0$ there is an L_r satisfying

$$\begin{aligned} \int |\widehat{H}_j^*(u)| du &= \int_{\{|u| \leq \lambda b_j\}} |\widehat{H}_j^*(u)| du + \int_{\{|u| \geq \lambda b_j\}} |\widehat{H}_j^*(u)| du \\ &\leq \int_{\{|u| \leq \lambda b_j\}} (Ae^{-d|u|^c} + L_r j^{-r}) du + B\rho^j \int g(u) du. \end{aligned}$$

for all $j \geq 2n_0$. Here the integrable function $g(u) \geq 0$ is the one occurring in the Statement (iii) of Lemma 16. Similarly, the second statement follows. \square

The next result deals with the situation of the second statement of Theorems 3.

Lemma 18. *Suppose that $\int |E[e^{iu\xi_1}]|^{n_0} du < \infty$ for some integer $n_0 > 0$. Then for all $j \geq n_0$ the conclusions of the preceding Lemma 17 hold for $\widehat{H}_j(u)$ itself.*

Proof. Because Statement (iii) of Lemma 16 holds for $\widehat{H}_j(u)$ itself for all $j \geq n_0$ with $g(v) = |E[e^{iv\xi_1}]|^{n_0}$, the proof follows as in the previous lemma using Lemma 12. \square

6 Proof of Proposition 6

We first prove the result under the conditions of Theorem 2 (*assumed to hold henceforth without further mentioning*), which proof will then essentially contain the proof under the conditions of Theorem 3. Throughout below the constants ν_n are such that

$$n^{-1}\nu_n \rightarrow 0 \text{ and } \nu_n^{-1}\beta_n \rightarrow 0.$$

Such a choice of ν_n is possible because $n^{-1}\beta_n \rightarrow 0$. Further, ρ will always stand for a constant with $0 < \rho < 1$. For the function $f_a(y) = f(ay)$, $a > 0$, the identities

$$\int M_{f_a, \frac{\eta}{a}}(y)dy = \frac{1}{a} \int M_{f, \eta}(y)dy, \quad \int m_{f_a, \frac{\eta}{a}}(y)dy = \frac{1}{a} \int m_{f, \eta}(y)dy, \quad (20)$$

which follow in view of $M_{f_a, \frac{\eta}{a}}(y) = M_{f, \eta}(ay)$ and $m_{f_a, \frac{\eta}{a}}(y) = m_{f, \eta}(ay)$, will be invoked repeatedly. In particular, because $\int M_{|f|, \eta}(y)dy < \infty$ for some $\eta > 0$,

$$\limsup_{n \rightarrow \infty} \int M_{|f_n|, \nu_n^{-1}}(y)dy = \limsup_{n \rightarrow \infty} \int M_{|f|, \beta_n \nu_n^{-1}}(y)dy \leq C. \quad (21)$$

We begin with following result, where and elsewhere **recall that** $f_n(y) = \beta_n f(\beta_n y)$.

Proposition 19.

$$\sup_x E[|f_n(\gamma_n^{-1}S_k + x)|] \leq C(\gamma_n \gamma_k^{-1} + \nu_n \rho^k) \int M_{|f|, \beta_n \nu_n^{-1}}(y)dy.$$

Proof. We have, with $g_n(j) = \gamma_n^{-1} \sum_{k=0}^j c_k$ (recall that $\gamma_n = n^{H-1/\alpha} R(n)b_n$)

$$\gamma_n^{-1}S_k = \sum_{j=-\infty}^0 (g_n(k-j) - g_n(1-j))\xi_j + \sum_{j=1}^k g_n(k-j)\xi_j = S_{nk}^{**} + \gamma_n^{-1}\gamma_k S_k^*$$

where $S_k^* = \sum_{j=1}^k g_k(k-j)\xi_j$. Note that S_k^* and S_{nk}^{**} are independent. Hence

$$\begin{aligned} E[|f_n(\gamma_n^{-1}S_k + x)|] &= E\left[\int |f_n(\gamma_n^{-1}\gamma_k y)| dH_k(y - \gamma_k^{-1}\gamma_n(x + S_{nk}^{**}))\right] \\ &= \int |f_{nk}| dH_{*k} \end{aligned}$$

where H_k is the distribution of S_k^* , $H_{*k}(y) = H_k(y - \gamma_k^{-1}\gamma_n(x + S_{n,k}^{**}))$ and $f_{nk}(y) = f_n(\gamma_n^{-1}\gamma_k y)$. Noting that S_k^* is in the form of (15), Lemma 14 (with the measure ν taken to be identically zero) gives $\int |f_{nk}| dH_{*k} \leq \int M_{|f_{nk}|, \nu_n^{-1}\gamma_n\gamma_k^{-1}} d(H_{*k} * K_{\nu_n^{-1}\gamma_n\gamma_k^{-1}})$. The r.h.s. is bounded by $\left(\int M_{|f_{nk}|, \nu_n^{-1}\gamma_n\gamma_k^{-1}}(y) dy\right) \left(\int \left|\widehat{H}_k(u)\right| \left|\widehat{K}_{\nu_n^{-1}\gamma_n\gamma_k^{-1}}(u)\right| du\right)$ because for a probability measure P with $\int \left|\widehat{P}(u)\right| du < \infty$, its density $p(y)$ satisfies

$$\sup_y p(y) \leq \int \left|\widehat{P}(u)\right| du. \quad (22)$$

($H_{*k} * K_\eta$ has a density function.) Now $\int \left|\widehat{H}_k(u)\widehat{K}_{\nu_n^{-1}\gamma_n\gamma_k^{-1}}(u)\right| du \leq C(1 + \rho^k \nu_n \gamma_n^{-1} \gamma_k)$, according to Lemma 15. Also $\int M_{|f_{nk}|, \nu_n^{-1}\gamma_n\gamma_k^{-1}}(y) dy = \gamma_n \gamma_k^{-1} \int M_{|f|, \beta_n \nu_n^{-1}}(y) dy$ in view of (20). Hence the proof follows. \square

Note that

$$\sup_{\varepsilon, x} E \left[\int_{-\infty}^{\infty} |f_n(\gamma_n^{-1} S_k + x + z\varepsilon)| \phi(z) dz \right] \leq \sup_x E [|f_n(\gamma_n^{-1} S_k + x)|] \quad (23)$$

Further $E [|f_n^2(\gamma_n^{-1} S_k + x)|] = \beta_n^2 E [|f^2(\beta_n(\gamma_n^{-1} S_k + x))|]$, and hence by Proposition 19

$$\sup_x E [|f_n^2(\gamma_n^{-1} S_k + x)|] \leq C \beta_n (\gamma_n \gamma_k^{-1} + \nu_n \rho^k) \int M_{f^2, \beta_n \nu_n^{-1}}(y) dy.$$

Now note that $n^{-1} \sum_{k=1}^n (\gamma_n \gamma_k^{-1} + \nu_n \rho^k) \leq C$ because $\gamma_k = k^H u(k)$ with $0 < H < 1$ and $u(k)$ slowly varying. Thus, by (20) and because $n^{-1} \beta_n \rightarrow 0$,

$$\left. \begin{aligned} & \sup_x \frac{1}{n^2} E \left[\sum_{k=1}^n |f_n^2(\gamma_n^{-1} S_k + x)| \right] \\ & \sup_{\varepsilon, x} \frac{1}{n^2} E \left[\sum_{k=1}^n \int_{-\infty}^{\infty} |f_n^2(\gamma_n^{-1} S_k + x + z\varepsilon)| \phi(z) dz \right] \end{aligned} \right\} \rightarrow 0 \quad (24)$$

Remark 7. If we let $f^{(\tau)}(y) = f(y) \mathbb{I}(|y| > \tau)$, then $\sup_x \beta_n E [|f^{(\tau)}(\beta_n(\gamma_n^{-1} S_k + x))|]$ is bounded by $C(\gamma_n \gamma_k^{-1} + \nu_n \rho^k) \int M_{|f^{(\tau)}|, \beta_n \nu_n^{-1}}(y) dy$, in view of Proposition 19, where $\lim_{\tau \rightarrow \infty} \limsup_{n \rightarrow \infty} \int M_{|f^{(\tau)}|, \beta_n \nu_n^{-1}}(y) dy = 0$ if $\int M_{|f|, \eta}(y) dy < \infty$ for some $\eta > 0$. Then without loss of generality $f(y)$ in Theorem 2 can be taken to have a compact support; in particular the integrability of $M_{f^2, \eta}(y)$ can be relaxed to local integrability.

The next Lemma will ease the computations to be carried out further on.

Lemma 20. *Let $Z_k, k \geq 1$, be i.i.d. standard Gaussian random variables independent of $\{\xi_j : -\infty < j < \infty\}$. Then for each $\varepsilon > 0$,*

$$\sup_{\varepsilon, t, x} E \left[\left\{ \frac{1}{n} \sum_{k=1}^{[nt]} \left(f_n(\gamma_n^{-1} S_k + x + Z_k \varepsilon) - \int_{-\infty}^{\infty} f_n(\gamma_n^{-1} S_k + x + z\varepsilon) \phi(z) dz \right) \right\}^2 \right] \rightarrow 0.$$

Proof. Define the σ -fields $F_j = \sigma(S_1, \dots, S_n, Z_1, \dots, Z_j)$. Because $\{Z_k\}$ and $\{S_k\}$ are independent, $E[f_n(\gamma_n^{-1}S_k + x + Z_k\varepsilon) | F_{k-1}] = \int_{-\infty}^{\infty} f_n(\gamma_n^{-1}S_k + x + z\varepsilon) \phi(z) dz$. The differences $\frac{1}{n} \{f_n(\gamma_n^{-1}S_k + x + Z_k\varepsilon) - E[f_n(\gamma_n^{-1}S_k + x + Z_k\varepsilon) | F_{k-1}]\}$ form martingale differences with respect to $\{F_k, k \geq 1\}$. Hence the expectation in the statement of the lemma is bounded by $E\left[\frac{1}{n^2} \sum_{k=1}^n \int_{-\infty}^{\infty} f_n^2(\gamma_n^{-1}S_k + x + z\varepsilon) \phi(z) dz\right]$. The proof follows by (24) \square

Final arguments of the Proof of Proposition 9. For notational convenience, and because all the bounds derived below will be independent of x , we restrict to the case $x = 0$. In the same way, we restrict to $t = 1$. In view of the preceding Lemma 20, it is enough to show that $E\left[\left\{\frac{1}{n} \sum_{k=1}^{\lfloor nt \rfloor} (f_n(\gamma_n^{-1}S_k) - f_n(\gamma_n^{-1}S_k + Z_k\varepsilon))\right\}^2\right] \rightarrow 0$ by letting first $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$. Then, in view of (24), it is enough to show that

$$\frac{1}{n^2} \sum_{j=1}^n \sum_{k=j+1}^n E[(f_n(\gamma_n^{-1}S_j) - f_n(\gamma_n^{-1}S_j + Z_j\varepsilon))(f_n(\gamma_n^{-1}S_k) - f_n(\gamma_n^{-1}S_k + Z_k\varepsilon))] \quad (25)$$

converges to 0 by first letting $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$. Recall that for $k > j$,

$$\gamma_n^{-1}S_k = \sum_{l=-\infty}^0 (g_n(k-l) - g_n(1-l))\xi_l + \sum_{l=1}^j g_n(k-l)\xi_l + \sum_{l=j+1}^k g_n(k-l)\xi_l$$

with $g_n(j) = \gamma_n^{-1} \sum_{k=0}^j c_k$ and $\gamma_n = n^{H-1/\alpha} R(n) b_n$. This is same as

$$\gamma_n^{-1}S_k = S_{nkj}^{**} + \gamma_n^{-1}\gamma_{k-j}S_{k-j}^*$$

where $S_{k-j}^* = \sum_{q=0}^{k-j-1} g_{k-j}(q)\xi_{k-q}$. Because S_{k-j}^* and S_{nkj}^{**} are independent,

$$\begin{aligned} & E[f_n(\gamma_n^{-1}S_j) f_n(\gamma_n^{-1}S_k + Z_k\varepsilon)] \\ &= E\left[f_n(\gamma_n^{-1}S_j) \int f_n(\gamma_n^{-1}\gamma_{k-j}y) d(H_{k-j} * G_{\varepsilon_{nkj}})(y - \gamma_{k-j}^{-1}\gamma_n S_{nkj}^{**})\right] \end{aligned}$$

where H_{k-j} is the distribution of S_{k-j}^* , G_σ is Gaussian with mean 0 and variance σ^2 and

$$\varepsilon_{nkj} = \varepsilon \gamma_{k-j}^{-1} \gamma_n.$$

Using similar identities, it follows that (25) can be written in the form

$$\frac{1}{n^2} \sum_{j=1}^n E\left[(f_n(\gamma_n^{-1}S_j) - f_n(\gamma_n^{-1}S_j + Z_j\varepsilon)) \sum_{k=j+1}^n \int f_{nkj} d(H_{k-j}^\# - H_{k-j}^\# * G_{\varepsilon_{nkj}})\right] \quad (26)$$

where we have set $f_{nkj}(y) = f_n(\gamma_n^{-1}\gamma_{k-j}y)$ and $H_{k-j}^\#(y) = H_{k-j}(y - \gamma_{k-j}^{-1}\gamma_n S_{nkj}^{**})$. We first show that, for every $\delta > 0$, the quantity

$$\frac{1}{n^2} \sum_{j=\lfloor n\delta \rfloor}^n E\left[(f_n(\gamma_n^{-1}S_j) - f_n(\gamma_n^{-1}S_j + Z_j\varepsilon)) \sum_{k=j+\lfloor j\delta \rfloor}^n \int f_{nkj} d(H_{k-j}^\# - H_{k-j}^\# * G_{\varepsilon_{nkj}})\right] \quad (27)$$

converges to 0, first by letting $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$. By taking $v = H_{k-j}^\#$, $\mu = H_{k-j}^\# * G_{\varepsilon_{nkj}}$ and $K_\eta = K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}$ in Lemma 14, one gets upper and lower bounds for

$$\int f_{nkj} d \left(H_{k-j}^\# * G_{\varepsilon_{nkj}} - H_{k-j}^\# \right). \quad (28)$$

By looking at these bounds, it is clear that we need to obtain bounds for the following

$$(c1) \quad \int M_{f_{nkj}, \nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} d \left(H_{k-j}^\# * G_{\varepsilon_{nkj}} * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} - H_{k-j}^\# * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right)$$

$$(c2) \quad \int \left(M_{f_{nkj}, 2\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} - f_{nkj} \right) dH_{k-j}^\#$$

$$(c3) \quad \int \left(f_{nkj} - m_{f_{nkj}, 2\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) dH_{k-j}^\#.$$

Regarding (c2) we have, again using Lemma 14 with $v = 0$ and $K_\eta = K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}$

$$\begin{aligned} & \int \left(M_{f_{nkj}, 2\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} - f_{nkj} \right) dH_{k-j}^\# \\ & \leq \int \left(M_{f_{nkj}, 3\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} - m_{f_{nkj}, \nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) d \left(H_{k-j}^\# * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) \\ & \leq \int \left(M_{f_{nkj}, 3\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}(y) - m_{f_{nkj}, 3\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}(y) \right) dy \int \left| \widehat{H}_{k-j}(u) \right| \left| \widehat{K}_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}(u) \right| du \\ & \leq C \gamma_n \gamma_{k-j}^{-1} \left(\int \left(M_{f_n, \nu_n^{-1}}(y) - m_{f_n, \nu_n^{-1}}(y) \right) dy \right) (1 + \rho^{k-j} \nu_n \gamma_n^{-1} \gamma_{k-j}) \\ & = C (\gamma_n \gamma_{k-j}^{-1} + \nu_n \rho^{k-j}) \int \left(M_{f, \beta_n \nu_n^{-1}}(y) - m_{f, \beta_n \nu_n^{-1}}(y) \right) dy \end{aligned} \quad (29)$$

where we have used (22), Statement (i) of Lemma 15 (S_{k-j}^* is in the form (15)) and the identity (20). In the same way, the final bound (29) holds for (c3) also. Now consider (c1).

Let $h_{k-j}(y)$ be the Lebesgue density of $H_{k-j}^\# * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}$. Then

$$\begin{aligned} & \int M_{f_{nkj}, \nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} d \left(H_{k-j}^\# * G_{\varepsilon_{nkj}} * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} - H_{k-j}^\# * K_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) \\ & = \int M_{f_{nkj}, \nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}}(y) \left[\int h_{k-j}(y - z\varepsilon_{nkj}) \phi(z) dz - h_{k-j}(y) \right] dy. \end{aligned} \quad (30)$$

Now (recall that $H_{k-j}^\#(y) = H_{k-j}(y - \gamma_{k-j}^{-1}\gamma_n S_{nkj}^{**})$)

$$\begin{aligned} & \left| \int h_{k-j}(y - z\varepsilon_{nkj}) \phi(z) dz - h_{k-j}(y) \right| \\ & = \left| \int \left(e^{-u^2 \varepsilon_{nkj}^2 / 2} - 1 \right) e^{iu(y - \gamma_{k-j}^{-1}\gamma_n S_{nkj}^{**})} \left(\widehat{H}_{k-j} * \widehat{K}_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) (u) du \right| \\ & \leq \int \left| e^{-u^2 \varepsilon_{nkj}^2 / 2} - 1 \right| \left| \left(\widehat{H}_{k-j} * \widehat{K}_{\nu_n^{-1}\gamma_n\gamma_{k-j}^{-1}} \right) (u) \right| du. \end{aligned}$$

Using this and the Statement (ii) of Lemma 15, (30) is bounded in absolute value by

$$\begin{aligned}
& \left(\int M_{|f_{nkj}|, \nu_n^{-1} \gamma_n \gamma_{k-j}^{-1}}(y) dy \right) \left(\int \left| e^{-u^2 \varepsilon_{nkj}^2 / 2} - 1 \right| \left| \left(\widehat{H}_{k-j} * \widehat{K}_{\nu_n^{-1} \gamma_n \gamma_{k-j}^{-1}} \right) (u) \right| du \right) \\
& \leq C \left(\varepsilon_{nkj}^2 + \rho^{k-j} \nu_n \gamma_n^{-1} \gamma_{k-j} \right) \gamma_n \gamma_{k-j}^{-1} \left(\int M_{|f|, \beta_n \nu_n^{-1}}(y) dy \right) \\
& \leq C \left(\varepsilon_{nkj}^2 \gamma_n \gamma_{k-j}^{-1} + \nu_n \rho^{k-j} \right). \tag{31}
\end{aligned}$$

Thus, summing (29) and (30), (28) is bounded in absolute value by a constant multiple of $(Q(n) + \varepsilon_{n,k-j}^2) \gamma_n \gamma_{k-j}^{-1} + (Q(n) + 1) \nu_n \rho^{k-j}$ where

$$Q(n) = \int (M_{f, \beta_n \nu_n^{-1}}(y) - m_{f, \beta_n \nu_n^{-1}}(y)) dy.$$

In addition, $E[|f_n(\gamma_n^{-1} S_j) - f_n(\gamma_n^{-1} S_j + Z_j \varepsilon)|] \leq C(\gamma_n \gamma_j^{-1} + \nu_n \rho^j)$ by Proposition 19 and (23). Thus (27) is bounded in absolute value by a constant multiple of

$$\frac{1}{n^2} \sum_{j=[n\delta]}^n (\gamma_n \gamma_j^{-1} + \nu_n \rho^j) \sum_{k=j+[j\delta]}^n ((Q(n) + \varepsilon_{nkj}^2) \gamma_n \gamma_{k-j}^{-1} + (Q(n) + 1) \nu_n \rho^{k-j}). \tag{32}$$

Now note that $\varepsilon_{nkj} = \varepsilon n^H (k-j)^{-H} u(n) (u(k-j))^{-1}$ where the slowly varying $u(n)$ is such that $\gamma_n = n^H u(n)$. When $k \geq j + [j\delta]$ and $j \geq [n\delta]$, there is a constant $C(\delta)$ depending on δ such that $n^H (k-j)^{-H} \leq C(\delta)$ for all sufficiently large n . Hence one can check that $\frac{1}{n^2} \sum_{j=[n\delta]}^n \gamma_n \gamma_j^{-1} \sum_{k=j+[j\delta]}^n \varepsilon_{n,k-j}^2 \gamma_n \gamma_{k-j}^{-1} \leq C_*(\delta) \varepsilon^2$ for some constant $C_*(\delta)$. In addition $\frac{1}{n^2} \sum_{j=1}^n \gamma_n \gamma_j^{-1} \sum_{k=j+1}^n \nu_n \rho^{k-j} \rightarrow 0$, $\frac{1}{n^2} \sum_{j=1}^n \nu_n \rho^j \sum_{k=j+1}^n \gamma_n \gamma_{k-j}^{-1} \rightarrow 0$ and $\frac{1}{n^2} \sum_{j=1}^n \nu_n \rho^j \sum_{k=j+1}^n \nu_n \rho^{k-j} \rightarrow 0$. (Recall that $n^{-1} \nu_n \rightarrow 0$.) Also $Q(n) \rightarrow 0$ by (7). Thus for each $\delta > 0$, (27) converges to 0 as $n \rightarrow \infty$ and $\varepsilon \rightarrow 0$. Hence it remains to show that the difference between (26) and (27) converges to 0, first by letting $n \rightarrow \infty$ and then $\delta \rightarrow 0$. The same arguments used in Proposition 19 shows that $\left| \int f_n(\gamma_n^{-1} \gamma_{k-j} y) d \left(H_{k-j}^\# * G_{\varepsilon_{nkj}} - H_{k-j}^\# \right) (y) \right|$ is bounded by $C(\gamma_n \gamma_{k-j}^{-1} + \nu_n \rho^{k-j})$. Hence, it can be seen that it is enough to show that $\frac{1}{n^2} \sum_{j=1}^{[n\delta]} \gamma_n \gamma_j^{-1} \sum_{k=j+1}^n \gamma_n \gamma_{k-j}^{-1}$ and $\frac{1}{n^2} \sum_{j=[n\delta]}^n \gamma_n \gamma_j^{-1} \sum_{k=j+1}^{j+[j\delta]} \gamma_n \gamma_{k-j}^{-1}$ converges to 0, first by letting $n \rightarrow \infty$ and then $\delta \rightarrow 0$. This is true for the first of these because it is bounded by $\frac{1}{n^2} \sum_{j=1}^{[n\delta]} \gamma_n \gamma_j^{-1} \sum_{l=1}^n \gamma_n \gamma_l^{-1}$. The same is true for the second one because it can be rewritten as

$$\begin{aligned}
\frac{1}{n^2} \sum_{j=[n\delta]}^n \gamma_n \gamma_j^{-1} \sum_{l=1}^{[j\delta]} \gamma_n \gamma_l^{-1} & \sim C \frac{\gamma_n^2}{n^2} \sum_{j=[n\delta]}^n \gamma_j^{-1} [j\delta] \gamma_{[j\delta]}^{-1} \\
& \sim C \frac{\delta^{1-H} \gamma_n^2}{n^2} \sum_{j=[n\delta]}^n j \gamma_j^{-2} \sim C \delta^{1-H}, \text{ as } n \rightarrow \infty
\end{aligned}$$

for each $\delta > 0$, where we have used the fact $j\gamma_j^{-2}$ is regularly varying with index $1 - 2H > -1$. This completes the proof of Proposition 6 under the assumptions of Theorems 2.

Now consider the proof under the assumptions of Theorems 3. First consider the first part of Theorem 3. Because $\sup_y |f(y)| \leq C$, we have $\frac{1}{n} \sum_{j=1}^{2n_0-1} |f_n(\gamma_n^{-1}S_j + x)| \leq \frac{Cn_0\beta_n}{n}$ where $\frac{\beta_n}{n} \rightarrow 0$. Hence one can restrict to the sum $\frac{1}{n} \sum_{j=2n_0}^n f_n(\gamma_n^{-1}S_j + x)$. Proceeding as in the proof of Proposition 19 and using Statement (i) of Lemma 16 and the first part of Lemma 17, for any $r > 0$ there is an L_r such that $E[|f_n(\gamma_n^{-1}S_k + x)|]$ is bounded by

$$\int |f_n(\gamma_n^{-1}\gamma_k y)| dy \left(\int |\widehat{H}_k^*(u)| du \right) + \frac{C\beta_n L_r}{k^r} \leq C\gamma_n\gamma_k^{-1} \int |f(y)| dy + \frac{C\beta_n L_r}{k^r}$$

for all $k \geq 2n_0$, where H_k^* is as in Lemma 16 corresponding to the distribution H_k of S_k^* . Hence, (24) (with the sums restricted to $2n_0 \leq k \leq n$) also holds, because $\int f^2(y) dy \leq (\sup_y |f(y)|) \int |f(y)| dy$. It remains to deal with (26), where as above, one can restrict the sum to j and k such that $k > j + 2n_0$ and $j \geq 2n_0$. In the same way as above but using the second part of Lemma 17, (28) is bounded by $C\gamma_n\gamma_{k-j}^{-1} (\varepsilon_{nkj}^2 + \rho^{k-j}) + \frac{C\beta_n L_r}{(k-j)^r}$ when $k > j + 2n_0$. Thus the same arguments used earlier under the assumptions of Theorem 2 become applicable when $r > 1$. (Now $\gamma_n\gamma_{k-j}^{-1}\rho^{k-j} + \frac{\beta_n L_r}{(k-j)^r}$ plays the role of the earlier $\nu_n\rho^{k-j}$ where only the facts $n^{-1}\nu_n \rightarrow 0$ and $\sum_{k=j+1}^n \rho^{k-j} < C$ were used.) This completes the proof of Proposition 6 under the first part of Theorem 3.

Regarding the proof under the second part of Theorem 3, note that we are considering the sum $\frac{1}{n} \sum_{j=n_0}^n f_n(\gamma_n^{-1}S_j + x)$ restricted to $n_0 \leq j \leq n$. Here we use the conclusions of Lemma 18, which hold for H_j itself for $j \geq n_0$. The bound in Proposition 19 now becomes $C\gamma_n\gamma_k^{-1} \int |f(y)| dy$ and the conclusion (24) (with the sums restricted to $n_0 \leq k \leq n$) also holds. Also, in dealing with (26) one can restrict the sum to j and k such that $k > j + n_0$ (and $j \geq n_0$), because

$$|E[f_n(\gamma_n^{-1}S_j + x) f_n(\gamma_n^{-1}S_k + x)]| \leq E^{\frac{1}{2}}[f_n^2(\gamma_n^{-1}S_j + x)] E^{\frac{1}{2}}[f_n^2(\gamma_n^{-1}S_k + x)]$$

so that, similar to (24), $\frac{1}{n^2} \sum_{j=n_0}^n |E[f_n(\gamma_n^{-1}S_j + x) f_n(\gamma_n^{-1}S_{j+q} + x)]| \rightarrow 0$ for each fixed $1 \leq q \leq n_0$. Further, as above, (28) is now bounded by $C\gamma_n\gamma_{k-j}^{-1} (\varepsilon_{nkj}^2 + \rho^{k-j})$ when $k > j + n_0$. Hence the proof of Proposition 6 is concluded. \square

Remark 8. Under the conditions of part two of Theorem 3, it is implicit in the preceding proof that, for every $0 < s_1 < s_2 \leq 1$, $\sup_x E \left[\left\{ \frac{1}{n} \sum_{j=[ns_1]}^{[ns_2]} f_n(\gamma_n^{-1}S_j + x) \right\}^2 \right]$ is bounded by $\frac{C}{n} \sum_{j=[ns_1]}^{[ns_2]} \gamma_n\gamma_j^{-1} \left(\frac{\beta_n}{n} \int f^2(y) dy + \frac{1}{n} \sum_{k=j+1}^{[ns_2]} \gamma_n\gamma_{k-j}^{-1} \right)$. Similar bounds hold under the first part of Theorem 3 and under Theorem 2. One can establish analogous bounds for $\sup_x E \left[\left\{ \frac{1}{n} \sum_{j=[ns_1]}^{[ns_2]} f_n(\gamma_n^{-1}S_j + x) \right\}^{2l} \right]$ for integers $l \geq 1$.

8 References

1. Akonom, J. (1993). Comportement asymptotique du temps d'occupation du processus des sommes partielles. *Ann. Inst. Henri. Poincaré.* 29, 57-81.
2. Astrauskas, A. (1983). Limit theorems for sums of linearly generated random variables. *Lith. Mat. J.* 23, 127-134.
3. Bhattacharya, R.N. and Ranga Rao, R. (1976). Normal approximation and asymptotic expansions. John Wiley, New York.
4. Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1987). Regular variation. Cambridge University Press, Cambridge.
5. Borodin, A.N. and Ibragimov, I.A. (1995). Limit theorems for functionals of random walks. *Proceedings of the Steklov Institute of Mathematics.* 195(2), 258+viii pages.
6. Gikhman, I.I. and Skorokhod, A.V. (1969). Introduction to the theory of random processes. Saunders, Philadelphia, PA. (Translated from the 1965 Russian Edition.)
7. Ibragimov, I.A. (1984). Some limit theorems for functionals of a random walk. *Dokl. Akad. Nauk. SSSR.* 276, 1049-1052. (English transl. in *Soviet Math. Dokl.* 29.)
8. Ibragimov, I.A. (1985). Théorèmes limites pour les marches aléatoires. *École d'Été de Probabilités de Saint Flour, XIII-1983. Lecture Notes in Math., Vol. 1117, 199-297.* Springer-Verlag, Berlin.
9. Kasahara, Y. and Maejima, M. (1988). Weighted sums of i.i.d. random variables attracted to integrals of stable processes. *Probab. Theory Related Fields.* 78, 75-96.
10. Kôno, N and Maejima, M. (1991). Self-similar stable processes with stationary increments. In *Stable processes and related topics*, Editors: Cambanis, S., Samorodnitsky, G. and Taqqu, M.S.
11. LeCam, L. (1960). Locally asymptotically normal families of distributions. *Univ. Calif. Publ. Statist.* 3, 27-98.
12. Maejima, A. (1989). Self-similar processes and limit theorems. *Sugaku Expositions.* 2, 103-123.

13. Park, J.Y. and Phillips, P.C.B. (1999). Asymptotics for nonlinear transformations of integrated time series. *Econometric Theory*. 15, 269-298.
14. Park, J.Y. and Phillips, P.C.B. (2001). Nonlinear regressions with integrated time series. *Econometrica*. 69, 117-161.
15. Prohorov, Yu.V. (1952). A local theorem for densities. *Dokl. Akad. Nauk. SSSR*. 83, 797-800.
16. Samorodnitsky, G. and Taqqu, M.S. (1994). *Stable non-Gaussian random processes: Stochastic models with infinite variance*. Chapman and Hall, New York.
17. Skorokhod, A.V. and Slobodenjuk, N.P. (1970). *Limit theorems for random walks*. "Naukova Dumka", Kiev. (Russian)
18. Tyurin, K. and Phillips, P.C.B. (1999). *The occupation density of fractional Brownian motion and some of its applications*. Working paper, Department of Economics, Yale University.