

ASYMPTOTIC EQUIVALENCE OF PROBABILISTIC SERIAL AND RANDOM PRIORITY MECHANISMS

YEON-KOO CHE AND FUHITO KOJIMA

Department of Economics, Columbia University
420 West 118th Street, 1016 IAB New York, NY 10027
yeonkooche@gmail.com

Department of Economics, Harvard University
1805 Cambridge St, Cambridge, MA 02138
fuhitokojima1979@gmail.com

ABSTRACT. The random priority (random serial dictatorship) mechanism is a common method for assigning objects to individuals. The mechanism is easy to implement and strategy-proof. However this mechanism is inefficient, as the agents may be made all better off by another mechanism that increases their chances of obtaining more preferred objects. Such an inefficiency is eliminated by the recent mechanism called probabilistic serial, but this mechanism is not strategy-proof. Thus, which mechanism to employ in practical applications has been an open question. This paper shows that these mechanisms become equivalent when the market becomes large. More specifically, given a set of object types, the random assignments in these mechanisms converge to each other as the number of copies of each object type approaches infinity. Thus, the inefficiency of the random priority mechanism becomes small in large markets. Our result gives some rationale for the common use of the random priority mechanism in practical problems such as student placement in public schools. *JEL Classification Numbers:* C70, D61, D63.

Keywords: random assignment, random priority, probabilistic serial, ordinal efficiency, asymptotic equivalence.

1. INTRODUCTION

Consider a mechanism designer's problem to assign a number of indivisible objects to agents who can consume at most one object each. University housing allocation and

Date: June 15, 2008.

We are grateful to Susan Athey, Mihai Manea, Al Roth, Parag Pathak, Michael Schwarz, Tayfun Sonmez, Rakesh Vohra, and seminar participants at Harvard Business School for helpful discussions. Yeon-Koo Che is grateful to the NSF Grant (SES#0721053) for financial support.

student placement in public schools are examples in real life.¹ A typical goal of the mechanism designer is to assign the objects efficiently and fairly, while eliciting the true preferences of the agents. The mechanism may need to satisfy other constraints as well. For example, considerations about fairness may prohibit monetary transfers and instead random assignments may be required. Also, the assignment often needs to be based on agents' reports of ordinal preferences over objects rather than full cardinal preferences, since elicitation of cardinal preferences may be difficult.² Two mechanisms have been regarded as promising solutions: the random priority mechanism (Abdulkadiroğlu and Sönmez 1998) and the probabilistic serial mechanism (Bogomolnaia and Moulin 2001).

In random priority, agents are ordered with equal probability and, for each realization of the ordering, the first agent in the ordering receives her most preferred object, the next agent receives his most preferred object among the remaining ones, and so on. Random priority is strategy-proof, that is, reporting ordinal preferences truthfully is a weakly dominant strategy for every agent. Moreover, random priority is ex-post efficient, that is, the assignment after the ordering lottery is resolved is Pareto efficient. This mechanism is simple and widely used in, among others, house allocation in universities and student placement in public schools. The procedure is also extended to various settings to accommodate constraints of specific applications, for example house allocation with existing tenants (Abdulkadiroğlu and Sönmez 1999) and student placement when some students have non-strict priorities for some schools (Abdulkadiroğlu, Pathak, and Roth 2005).

Although the random priority mechanism is ex-post efficient, it may result in unambiguous efficiency loss ex ante. Bogomolnaia and Moulin (2001) provide an example in which the random priority assignment is dominated by another random assignment that improves the chance of obtaining a more preferred object for each agent, in the sense of first-order stochastic dominance. Bogomolnaia and Moulin introduce the ordinal efficiency concept: a random assignment is ordinally efficient if it is not first-order stochastically dominated for all agents by any other random assignment. Ordinal efficiency is perhaps the most relevant efficiency concept in the context of assignment mechanisms based solely on ordinal preferences.

¹See Abdulkadiroğlu and Sönmez (1999) and Chen and Sönmez (2002) for application to house allocation, and Balinski and Sönmez (1999) and Abdulkadiroğlu and Sönmez (2003b) for student placement. For the latter application, Abdulkadiroğlu, Pathak, and Roth (2005) and Abdulkadiroğlu, Pathak, Roth, and Sönmez (2005) discuss practical considerations in designing student placement mechanisms in New York City and Boston.

²The pseudo-market mechanism proposed by Hylland and Zeckhauser (1979) is one of the few mechanisms proposed in the literature in which agents report their cardinal preferences over objects.

Bogomolnaia and Moulin propose the probabilistic serial mechanism as an alternative to the random priority mechanism. The basic idea is to regard each object as a continuum of “probability shares.” Each agent “eats” her most preferred available object (in probability share) with speed one at every point in time between 0 and 1. The probabilistic serial random assignment is defined as the profile of shares of objects eaten by agents by time 1. The probabilistic serial random assignment is ordinally efficient if all the agents report their ordinal preferences truthfully.

However, the probabilistic serial mechanism is not strategy-proof. In other words, an agent may receive a more desirable random assignment (with respect to her true expected utility function) by misreporting her ordinal preferences. Thus it has been unclear which of these two mechanisms, random priority or probabilistic serial, performs well in applications.

This paper shows that the random priority mechanism and the probabilistic serial mechanism become equivalent when the market becomes large. More specifically, given a set of object types, the random assignments in these mechanisms converge to each other as the number of copies of each object type approaches infinity. Thus inefficiency of the random priority mechanism becomes small in large markets. If the random priority mechanism is easy to implement, then our result gives some supports to the common use of the random priority mechanism as the market design in practical problems such as student placement in public schools.

In our model, the large market assumption means that there exist a large number of copies of each object type. This model includes several interesting cases. For instance, the replica economies model is a special case, as the number of copies of each object type is large in an economy that is replicated a large number of times. Also, the assumption is natural in several practical applications. In student placement in public schools, there are typically a large number of identical seats at each school. In the context of university housing allocation, the set of rooms may be partitioned into a number of categories by building and size, and all rooms of the same type may be treated to be identical.³ Our model may be applicable to these markets.

This paper investigates several additional issues as well. First, we define random priority and probabilistic serial directly in economies with continuum of agents and objects. We show that random priority and the probabilistic serial mechanisms in finite economies converge to those in the continuum economy. In that sense, we provide foundation of a

³For example, the assignment of graduate housing at Harvard University is based on the preferences of each student over eight types of rooms: two possible sizes (large and small) and four buildings.

modeling approach that directly studies economies with continuum of objects and agents. Second, we show that our equivalence is tight in the sense that in any finite economy, random priority and probabilistic serial can be different. We also present several extensions such as cases with existing priority and multi-unit demands.

Related literature. Pathak (2006) compares random priority and probabilistic serial using data in the assignment of about 8,000 students in the public school system of New York City. He finds that many students obtain a better random assignment in the probabilistic serial mechanism, but he notes that the difference seems small.

Kojima and Manea (2008) consider a large market that is similar to the current setting and find that truth-telling becomes a dominant strategy under the probabilistic serial mechanism when there are a large number of copies of each object type. While logically unrelated, our analysis provides intuition for their result. To see this point, first recall that truth-telling is a dominant strategy in the random priority mechanism. Since our result shows that the probabilistic serial mechanism is close to the random priority mechanism in a large economy, this observation implies that it is difficult to profitably manipulate the probabilistic serial mechanism. We note that, however, the result of Kojima and Manea (2008) cannot be derived from the current paper. This is because their assumptions are slightly different from ours and they establish a dominant strategy result in a large but finite economies, while our equivalence result holds only in the limit as the market size approaches infinity.

Manea (2006a) shows that random priority results in an ordinally inefficient assignment for most preference profiles when there are a large number of object types and the number of copies of each object type remains one. We note that his result does not contradict ours because of a number of differences. First, Manea (2006a) keeps the number of copies of each object type constant (at one) and increases the number of object types while our model increases the number of copies while keeping the number of object types fixed. Second, his theorem focuses on whether there is *some* ordinal inefficiency in the random priority assignment, while we investigate *how much* difference there is between the random priority and the probabilistic serial mechanisms, and how they change as the market size grows.

While analysis of large markets is relatively new in matching and resource allocation problems, it has a long tradition in many areas of economics. For example, in a pure exchange economy, the classical result of Hurwicz (1972) implies that the Walrasian mechanism is not strategy-proof. However Roberts and Postlewaite (1976) show that, under some conditions, the Walrasian mechanism becomes increasingly difficult to manipulate as

the market becomes large.⁴ Similarly, incentive properties of a large class of double auction mechanisms are studied by, among others, Gresik and Satterthwaite (1989), Rustichini, Satterthwaite, and Williams (1994), and Cripps and Swinkels (2006) Two-sided matching is an area closely related to our model. In that context, Roth and Peranson (1999), Immorlica and Mahdian (2005) and Kojima and Pathak (2008) show that the deferred acceptance algorithm proposed by Gale and Shapley (1962) becomes increasingly hard to manipulate as the number of participants becomes large. One of the notable features of the current paper is that we show the equivalence of apparently different mechanisms, beyond showing particular properties (such as incentive compatibility and efficiency) of given mechanisms.

Finally, there is a growing literature on random assignment and ordinal efficiency. Several characterizations of ordinal efficiency are given by Abdulkadiroğlu and Sönmez (2003a) and McLennan (2002).⁵ Kesten (2008) introduces two mechanisms, one of which is motivated by the random priority mechanism, and shows that these mechanisms are equivalent to the probabilistic serial mechanism. The probabilistic serial mechanism is generalized to cases with indifferences in preferences, existing property rights, and assignment of multiple objects to each agent by Katta and Sethuraman (2006), Yilmaz (2006), and Kojima (2008), respectively. In the scheduling problem (a special case of the current environment), Crès and Moulin (2001) show that the probabilistic serial mechanism is group strategy-proof and stochastically dominates the random priority mechanism but these two mechanisms converge to each other as the market size approaches infinity, and Bogomolnaia and Moulin (2002) give two characterizations of the probabilistic serial mechanism.

The rest of the paper proceeds as follows. Section 2 introduces the model. Section 3 defines the random priority mechanism and the probabilistic serial mechanism. Section 4 presents the main result. Section 5 provides discussion and concluding remarks. Proofs are found in the Appendix unless stated otherwise.

2. MODEL

For each $q \in \mathbb{N}$, consider a q -**economy**, $\Gamma^q = (N^q, (\pi_i)_{i \in N^q}, O)$, where N^q represents the set of agents and O represents the set of **proper object types** (we assume that O is identical for all q). There are $|O| = n$ object types, and each object type $a \in O$ has q copies available. There exist an infinite number of copies of a **null object** \emptyset , which is not

⁴See also Jackson (1992) and Jackson and Manelli (1997).

⁵A short constructive proof of the characterization by McLennan (2002) is given by Manea (2006b).

included in O . Let $\tilde{O} := O \cup \{\emptyset\}$. Each agent $i \in N$ has a **strict preference** represented by a permutation $\pi_i \in \Pi$ of \tilde{O} , where a given permutation $\pi_i : \{1, \dots, n+1\} \mapsto \tilde{O}$ lists for its j -th element $\pi_i(j)$ the agent's j -th most preferred object. (That is, agent i prefers a over b if and only if $\pi_i^{-1}(a) < \pi_i^{-1}(b)$.) For preference type π and for any $O' \subset \tilde{O}$,

$$Ch_\pi(O') := \{a \in O' \mid \pi^{-1}(a) \leq \pi^{-1}(b) \ \forall b \in O'\},$$

is the object that an agent of preference type π chooses if the set O' of objects are available to her.

The agents are partitioned into different preference types: $N^q = \{N_\pi^q\}_{\pi \in \Pi}$, where N_π^q is the set of the agents with preference $\pi \in \Pi$ in the q -economy. Let $m_\pi^q := \frac{|N_\pi^q|}{q}$ be the per-unit number of agents of type π in the q -economy.⁶ We assume, for each $\pi \in \Pi$, there exists $m_\pi^\infty \in \mathbb{R}_+$ such that $m_\pi^q \rightarrow m_\pi^\infty$ as $q \rightarrow \infty$. For $q \in \mathbb{N} \cup \{\infty\}$, let $m^q := \{m_\pi^q\}_{\pi \in \Pi}$. For any $q \in \mathbb{N} \cup \{\infty\}$, $O' \subset O$ and $a \in O'$, let

$$m_a^q(O') = \sum_{\pi \in \Pi} \{m_\pi^q \mid a \in Ch_\pi(O')\},$$

be the per unit number of agents whose most preferred object in O' is a in the q -economy. Throughout, we do not impose any restriction on the way in which the q -economy, Γ^q , grows with q (except that we assume the existence of the limit $m_\pi^\infty = \lim_{q \rightarrow \infty} m_\pi^q$ for each $\pi \in \Pi$).

A special case of interest is when the economy grows at a constant rate with q . We say that the family $\{\Gamma^q\}_{q \in \mathbb{N}}$ are **replica economies** if $m_\pi^q = m_\pi^\infty$ (or equivalently, $|N_\pi^q| = q|N_\pi^1|$) for all $q \in \mathbb{N}$ for all $\pi \in \Pi$, and call Γ^1 a **base economy** and Γ^q its **q -fold replica**.

Throughout, we focus on a symmetric random assignment in which all the agents with the same preference type π receive the same lottery over the goods. Formally, a **symmetric random assignment in the q -economy** is a mapping $\phi^q : \Pi \mapsto \Delta\tilde{O}$, where $\Delta\tilde{O}$ is the set of probability distributions over \tilde{O} , that satisfies the feasibility constraint $\sum_{\pi \in \Pi} \phi_a^q(\pi) \cdot |N_\pi^q| \leq q$, for each $a \in O$, where $\phi_a^q(\pi)$ represents the probability that a type π -agent receives the good a .⁷

It is useful to describe the limit economy (∞ -economy) independently. For this purpose, we assume that there exists a continuum of copies of objects in O and agents in N^∞ . More precisely, there exists a unit mass of each good in O , and the set of agent types Π is then endowed with a measure $\mu : \Pi \mapsto \mathbb{R}_+$ such that $\mu(\pi) = m_\pi^\infty$. **A symmetric**

⁶Given a set X , we denote the cardinality of X by $|X|$ or $\#X$.

⁷The symmetry assumption that all the agents with the same preference type π receive the same lottery is often called the ‘‘equal treatment of equals’’ axiom.

random assignment in the limit economy is then defined as $\phi^\infty : \Pi \mapsto \Delta \tilde{O}$ such that $\sum_{\pi \in \Pi} \phi_a^\infty(\pi) \cdot m_\pi^\infty \leq 1$ for each $a \in O$.

2.1. Ordinal Efficiency. Consider a q -economy (where $q \in \mathbb{N} \cup \{\infty\}$). A symmetric random assignment ϕ^q **ordinally dominates** another random assignment $\hat{\phi}^q$ **at** m^q if for each preference type π with $m_\pi^q > 0$ the lottery $\phi^q(\pi)$ first-order stochastically dominates the lottery $\hat{\phi}^q(\pi)$,

$$(2.1) \quad \sum_{\pi^{-1}(b) \leq \pi^{-1}(a)} \phi_b^q(\pi) \geq \sum_{\pi^{-1}(b) \leq \pi^{-1}(a)} \hat{\phi}_b^q(\pi) \quad \forall \pi, m_\pi^q > 0, \forall a \in \tilde{O},$$

with strict inequality for some π, a . The random assignment ϕ^q is **ordinally efficient at** m^q if it is not ordinally dominated at m^q by any other random assignment. If ϕ^q ordinally dominates $\hat{\phi}^q$ at m^q , then every agent of every preference type prefers their assignment under ϕ^q to the one under $\hat{\phi}^q$ according to any expected utility function with utility index consistent with their ordinal preferences.

We say that ϕ^q is **non-wasteful at** m^q if there exists no preference type $\pi \in \Pi$ with $m_\pi^q > 0$ and objects $a, b \in \tilde{O}$ such that $\pi^{-1}(a) < \pi^{-1}(b)$, $\phi_b^q(\pi) > 0$ and $\sum_{\pi' \in \Pi} \phi_a^q(\pi') m_{\pi'}^q < 1$.

Consider the binary relation $\triangleright(\phi^q, m^q)$ on \tilde{O} defined by

$$(2.2) \quad a \triangleright (\phi^q, m^q) b \iff \exists \pi \in \Pi, m_\pi^q > 0, \pi^{-1}(a) < \pi^{-1}(b) \text{ and } \phi_b^q(\pi) > 0.$$

In a setting in which each object has quota 1 and there exist an equal number of agents and objects, Bogomolnaia and Moulin show the equivalence of ordinal efficiency and acyclicity of this binary relation. Their characterization extends straightforwardly to our setting as follows (the proof is omitted).

Proposition 1. The random assignment ϕ^q is ordinally efficient at m^q if and only if the relation $\triangleright(\phi^q, m^q)$ is acyclic and ϕ^q is non-wasteful at m^q .

3. TWO COMPETING MECHANISMS

3.1. Random Priority Mechanism. We introduce the **random priority mechanism** (Bogomolnaia and Moulin 2001), also called the **random serial dictatorship** (Abulkadiroğlu and Sönmez 1998), which is widely used in practice. Given preferences of all the agents, the agents are ordered randomly, and each agent selects, according to the order, the most preferred object among the remaining ones. For our purpose, it is useful to model the random ordering procedure as follows: First, each agent i randomly draws a number f_i independently from a uniform distribution on $[0, 1]$. Second, the agent with the smallest draw receives her most preferred object, the agent with the second-smallest draw

receives his most preferred object from the remaining ones, and so forth (it suffices to only consider cases in which $f_i \neq f_j$ for any $i \neq j$, since $f_i = f_j$ occurs with probability zero). This procedure induces a random assignment. We let RP^q be the random assignment under the random priority mechanism in Γ^q .

The random assignment RP^q is characterized as follows. Fix an agent i of arbitrary preference π , and fix the draws $f_{-i} = (f_j)_{j \in N \setminus \{i\}} \in [0, 1]^{(|N^q|-1)}$ for all agents other than i . We then ask how low agent i 's draw should be for her to obtain a given object $a \in \tilde{O}$. Specifically, we characterize the **cutoff** $\hat{T}_a^q \in [0, 1]$ for each good $a \in O$, which represents *the largest value of draw that would allow agent i to claim a* . It is the critical value in $[0, 1]$ such that agent i can obtain a if and only if she draws f_i less than that value. The cutoffs depend on the random draws f_{-i} , so they are random. It is useful to characterize the random assignment RP^q through the cutoffs.

To begin, let $\hat{m}_{\pi'}^q(t, t') := \frac{\#\{i \in N^q \setminus \{i\} | f_i \in (t, t']\}}{q}$ denote the per-unit number of agents of type π' (except i if $\pi' = \pi$) who have draws in $(t, t']$. For any $O' \subset O$ and $a \in O'$, we let

$$\hat{m}_a^q(O'; t, t') = \sum_{\pi' \in \Pi} \{\hat{m}_{\pi'}^q(t, t') | a \in Ch_{\pi'}(O')\},$$

be the per-unit number of agents in $N^q \setminus \{i\}$ whose most preferred object in O' is a and who have draws in $(t, t']$.

We then characterize the cutoffs for i by the following sequence of steps. Let $\hat{O}^q(0) = \tilde{O}$, $\hat{t}^q(0) = 0$, and $\hat{x}_a^q(0) = 0$ for every $a \in \tilde{O}$. Given $\hat{O}^q(0), \hat{t}^q(0), \{\hat{x}_a^q(0)\}_{a \in \tilde{O}}, \dots, \hat{O}^q(v-1), \hat{t}^q(v-1), \{\hat{x}_a^q(v-1)\}_{a \in \tilde{O}}$, we let $\hat{t}_\emptyset^q := 1$ and for each $a \in O$, define

$$(3.1) \quad \hat{t}_a^q(v) = \sup \left\{ t \in [0, 1] \mid \hat{x}_a^q(v-1) + \hat{m}_a^q(\hat{O}^q(v-1); \hat{t}^q(v-1), t) < 1 \right\},$$

$$(3.2) \quad \hat{t}^q(v) = \min_{a \in \hat{O}(v-1)} \hat{t}_a^q(v),$$

$$(3.3) \quad \hat{O}^q(v) = \hat{O}^q(v-1) \setminus \{a \in \hat{O}^q(v-1) | \hat{t}_a^q(v) = \hat{t}^q(v)\},$$

$$(3.4) \quad \hat{x}_a^q(v) = \hat{x}_a^q(v-1) + \hat{m}_a^q(\hat{O}^q(v-1); \hat{t}^q(v-1), \hat{t}^q(v)).$$

The last step $\hat{v}^q := \min\{v' | \hat{t}^q(v') = 1\}$ is well defined since O is finite. For each $a \in O$, its cutoff is given by $\hat{T}_a^q := \{\hat{t}^q(v) | \hat{t}_a^q(v) = \hat{t}^q(v)\}$ if the set is nonempty, or else $\hat{T}_a^q = 1$.

This characterization is explained as follows. Each step determines the cutoff of a good. Suppose steps 1 through $v-1$ have determined the $v-1$ cutoffs for $v-1$ objects. In particular, by the end of step $v-1$, agents with draws less than $\hat{t}^q(v-1)$ have consumed entire q copies of these goods and a fraction $\hat{x}_b^q(v-1)$ of each remaining good $b \in O^q(v-1)$.

Suppose the good $a \in \hat{O}^q(v-1)$ is next to be consumed away, by agents with draws less than its cutoff, $\hat{t}^q(v) = \hat{T}_a^q$. An agent with draw $f \in (\hat{t}^q(v-1), \hat{T}_a^q]$ will consume the good if

and only if she prefers a to all other remaining goods. The total number of all such agents is $q \cdot \hat{m}_a^q(\hat{O}^q(v-1); \hat{t}^q(v-1), \hat{T}_a^q)$, and they consume a fraction $\hat{m}_a^q(\hat{O}^q(v-1); \hat{t}^q(v-1), \hat{T}_a^q)$ of that good. Hence, the total fraction of a consumed by all agents with draws less than \hat{T}_a^q must be

$$\hat{x}_a^q(v-1) + \hat{m}_a^q(\hat{O}^q(v-1); \hat{t}^q(v-1), \hat{T}_a^q).$$

For \hat{T}_a^q to be the cutoff for a , this fraction must be no greater than one and must equal one if $\hat{T}_a^q < 1$. This condition requires \hat{T}_a^q to equal $\hat{t}_a^q(v)$, defined in (3.1). That a is the first to be consumed away among the remaining objects is (3.2). The last two equations reset the remaining set of goods and the fractions consumed by step v , thus continuing on the recursive procedure.

For each a , let

$$\hat{\tau}_a^q(\pi) := \min\{\hat{t}^q(v) \leq \hat{T}_a^q | a \in Ch_\pi(\hat{O}^q(v-1))\}$$

be the minimum value of draw for an agent with preference π , to choose a (again if the minimum is well defined, or else let $\hat{\tau}_a^q(\pi) := \hat{T}_a^q$). For instance, if all goods that agent $i \in N_\pi^q$ prefers to good a have lower cutoffs than a and b has the highest cutoff among those preferred to a , then $\hat{\tau}_a^q(\pi) = \hat{T}_b^q$. In that case, agent i will obtain a if and only if her draw f_i lies between two cutoffs \hat{T}_b^q and \hat{T}_a^q , as is depicted in Figure 1.




Figure 1: Cutoffs of objects under RP.

Therefore, the **random priority random assignment** is defined, for $i \in N_\pi^q$ and $a \in O$, as $RP_a^q(\pi) := \mathbb{E}[\hat{T}_a^q - \hat{\tau}_a^q(\pi)]$, where the expectation \mathbb{E} is taken with respect to $f_{-i} = (f_j)_{j \neq i}$ which is distributed i.i.d uniformly on $[0, 1]$.

The random priority mechanism is widely used in practice. Moreover, the mechanism is **strategy-proof**, that is, reporting the true ordinal preferences is a dominant strategy for each agent. Moreover, it is **ex post efficient**, that is, the assignment after random draws are realized is Pareto efficient. However, the mechanism may result in an ordinaly inefficient allocation, as shown by the following example adapted from Bogomolnaia and Moulin (2001).

Example 1. Consider an economy Γ^1 with 2 types of proper objects, a and b , each having one copy. Let $N^1 = N_\pi^1 \cup N_{\pi'}^1$ be the set of agents, with $|N_\pi^1| = |N_{\pi'}^1| = 2$. Preferences of

the agents are specified by

$$\begin{aligned}(\pi(1), \pi(2), \pi(3)) &= (a, b, \emptyset), \\(\pi'(1), \pi'(2), \pi'(3)) &= (b, a, \emptyset).\end{aligned}$$

In this economy, the random assignments under RP^1 can be easily calculated to be

$$\begin{aligned}RP^1(\pi) &= (RP_a^1(\pi), RP_b^1(\pi), RP_\emptyset^1(\pi)) = \left(\frac{5}{12}, \frac{1}{12}, \frac{1}{2}\right), \\RP^1(\pi') &= (RP_a^1(\pi'), RP_b^1(\pi'), RP_\emptyset^1(\pi')) = \left(\frac{1}{12}, \frac{5}{12}, \frac{1}{2}\right).\end{aligned}$$

Each agent ends up with her less preferred object with positive probability, since two agents of any given preference type may get the two best draws, in which case the agent with the second best draw will take the less preferred object.⁸ Obviously, any two agents of different preferences can benefit from trading off the probability share of the less preferred object with that of the most preferred. In other words, the RP assignment is ordinally dominated by

$$\begin{aligned}\phi^1(\pi) &= \left(\frac{1}{2}, 0, \frac{1}{2}\right), \\ \phi^1(\pi') &= \left(0, \frac{1}{2}, \frac{1}{2}\right).\end{aligned}$$

Therefore the random priority assignment RP^1 is ordinally inefficient in this market.

Ordinal inefficiency of RP can be traced to the random and personalized features of its cutoffs. In Example 1, $\hat{T}_a^1 < \hat{T}_b^1$ occurs to agents in N_π^1 with positive probability, and $\hat{T}_a^1 > \hat{T}_b^1$ occurs to agents in $N_{\pi'}^1$ with positive probability. In the former case, an agent in N_π^1 may get b even though she prefers a to b . In the latter case, an agent in $N_{\pi'}^1$ may get a even though she prefers b to a . Hence both $a \triangleright (RP^1, m^1)b$ and $b \triangleright (RP^1, m^1)a$ occur, resulting in cyclicity of the relationship $\triangleright (RP^1, m^1)$. As we will see below, the probabilistic serial mechanism will admit a deterministic cutoffs that are common to all agents, and this feature ensures acyclicity of the binary relation \triangleright .

3.2. Probabilistic Serial Mechanism. Now we introduce the **probabilistic serial** mechanism, which is an adaptation of the mechanism proposed by Bogomolnaia and Moulin to our setting. The idea is to regard each object as a divisible good of “probability shares.” Each agent “eats” the best available object with speed one at every time $t \in [0, 1]$ (object a is available at time t if less than q share of a has been eaten away by time t).

⁸For instance, let $N_\pi^1 = \{1, 2\}$ and $N_{\pi'}^1 = \{3, 4\}$. The draws can be $f_1 < f_2 < f_3 < f_4$, in which case 1 gets a and 2 gets b and 3 and 4 get nothing.

The resulting profile of shares of objects eaten by agents by time 1 obviously corresponds to a symmetric random assignment, which we call the **probabilistic serial random assignment**.

Formally, the **symmetric simultaneous eating algorithm**,⁹ used to determine the probabilistic serial random assignment, is defined as follows.

PS mechanism in the finite economy. For q -economy, Γ^q , the assignment under the probabilistic serial mechanism is defined by the following sequence of steps. Let $O^q(0) = \tilde{O}$, $t^q(0) = 0$, and $x_a^q(0) = 0$ for every $a \in \tilde{O}$. Given $O^q(0), t^q(0), \{x_a^q(0)\}_{a \in \tilde{O}}, \dots, O^q(v-1), t^q(v-1), \{x_a^q(v-1)\}_{a \in \tilde{O}}$, we let $t_\emptyset^q := 1$ and for each $a \in O$, define

$$(3.5) \quad t_a^q(v) = \sup \{t \in [0, 1] \mid x_a^q(v-1) + m_a^q(O^q(v-1))(t - t^q(v-1)) < 1\},$$

$$(3.6) \quad t^q(v) = \min_{a \in O(v-1)} t_a^q(v),$$

$$(3.7) \quad O^q(v) = O^q(v-1) \setminus \{a \in O^q(v-1) \mid t_a^q(v) = t^q(v)\},$$

$$(3.8) \quad x_a^q(v) = x_a^q(v-1) + m_a^q(O^q(v-1))(t^q(v) - t^q(v-1)).$$

The last step $\bar{v}^q := \min\{v' \mid t^q(v') = 1\}$ is again well defined since O is finite. For each $a \in \tilde{O}$, define its **expiration date**: $T_a^q := \{t^q(v) \mid t_a^q(v) = t^q(v)\}$. The expiration date for good a is the time at which the eating of a is complete. When an agent starts eating a given good a depends on his preference. Note that, unlike the cutoffs in the random priority rule, the expiration dates are deterministic and common to all agents. Aside from this important difference, though, expiration dates in PS play a similar role as cutoffs in RP. In particular, they completely pin down the random assignment for the agents. To begin, for $\pi \in \Pi$, we let

$$\tau_a^q(\pi) := \min\{t^q(v) \leq T_a^q \mid a \in Ch_\pi(O^q(v-1))\}$$

if the minimum is well defined, or else let $\tau_a^q(\pi) := T_a^q$. Then, agent i 's probability of getting assigned to $a \in \tilde{O}$ is simply its duration of consumption of its preference type; i.e., $PS_a^q(\pi) = T_a^q - \tau_a^q(\pi)$ if $i \in N_\pi^q$.

PS mechanism in the limit economy. Although our primary interest is in a large but finite economy, it is useful to define the PS mechanism in the limit economy, for it will act as a benchmark for subsequent analysis. We again do so recursively. Let $O^\infty(0) = \tilde{O}$, $t^\infty(0) = 0$, and $x_a^\infty(0) = 0$ for every $a \in \tilde{O}$. Given $O^\infty(0), t^\infty(0), \{x_a^\infty(0)\}_{a \in \tilde{O}}, \dots,$

⁹Bogomolnaia and Moulin (2001) consider a broader class of simultaneous eating algorithms, where eating speeds may vary across agents and time.

$O^\infty(v-1), t^\infty(v-1), \{x_a^\infty(v-1)\}_{a \in \bar{O}}$, we let $t_\emptyset^\infty := 1$ and for each $a \in O$, define

$$(3.9) \quad t_a^\infty(v) = \sup \{t \in [0, 1] \mid x_a^\infty(v-1) + m_a^\infty(O^\infty(v-1))(t - t^\infty(v-1)) < 1\},$$

$$(3.10) \quad t^\infty(v) = \min_{a \in O^\infty(v-1)} t_a^\infty(v),$$

$$(3.11) \quad O^\infty(v) = O^\infty(v-1) \setminus \{a \in O^\infty(v-1) \mid t_a^\infty(v) = t^\infty(v)\},$$

$$(3.12) \quad x_a^\infty(v) = x_a^\infty(v-1) + m_a^\infty(O^\infty(v-1))(t^\infty(v) - t^\infty(v-1)).$$

Let \bar{v}^∞ such that $t^\infty(\bar{v}^\infty) = 1$. Consider the associated expiration dates: For each $a \in O$, $T_a^\infty := \{t^\infty(v) \mid t^\infty(v) = t_a^\infty(v)\}$ if the set is nonempty, or else $T_a^\infty := 1$. Likewise, the starting time for a for π is defined as

$$\tau_a^\infty(\pi) := \min\{t^\infty(v) \leq T_a^\infty \mid a \in Ch_\pi(O^q(v-1))\}$$

if the minimum is well defined, or else let $\tau_a^\infty(\pi) := T_a^\infty$. The **PS random assignment in limit** is then defined to be duration of eating each good: for $a \in O$, $PS_a^\infty(\pi) := T_a^\infty - \tau_a^\infty(\pi)$.

Adapting the argument of Bogomolnaia and Moulin (2001), we can show the following (the proof is omitted).

Proposition 2. For any $q \in \mathbb{N} \cup \{\infty\}$, PS^q is ordinally efficient.

Example 2. Consider replica economies $\{\Gamma^q\}_{q \in \mathbb{N}}$ with 2 types of proper objects, a and b , each having quota q in the q -fold replica. Let $N^q = N_\pi^q \cup N_{\pi'}^q$ be the set of agents in the q -fold replica, with N_π^q and $N_{\pi'}^q$ containing $2q$ agents each. Assume that the preferences of the agents are specified by

$$\begin{aligned} (\pi(1), \pi(2), \pi(3)) &= (a, b, \emptyset), \\ (\pi'(1), \pi'(2), \pi'(3)) &= (b, a, \emptyset). \end{aligned}$$

Note that Γ^1 corresponds to the market in Example 1.

For any $q \in \mathbb{N}$, the random assignments under PS^q can be easily calculated to be

$$\begin{aligned} PS^q(\pi) &= (PS_a^q(\pi), PS_b^q(\pi), PS_\emptyset^q(\pi)) = \left(\frac{1}{2}, 0, \frac{1}{2}\right), \\ PS^q(\pi') &= (PS_a^q(\pi'), PS_b^q(\pi'), PS_\emptyset^q(\pi')) = \left(0, \frac{1}{2}, \frac{1}{2}\right), \end{aligned}$$

which is ordinally efficient.

Unlike cutoffs in the random priority mechanism, the expiration dates in the probabilistic serial mechanism are deterministic and common to all agents. This explains, for

instance, why we cannot have a cycle on a binary relation $\triangleright(RP^q, m^q)$. To see this, suppose $a \triangleright (RP^q, m^q)b$. Then, there must be an agent who prefers a to b but ends up with b with positive probability. This is possible only if $T_a > T_b$; or else, by the time the agent finishes “eating” a (or something even better than a), b will have been completely eaten away. Based on this logic, a cycle on $\triangleright(RP^q, m^q)$ will require the order on the expiration dates to be cyclic. And this is impossible.

One main drawback of the probabilistic serial mechanism, as identified by Bogomolnaia and Moulin (2001), is that the mechanism is not strategy-proof. In other words, an agent may be made better off by reporting a false ordinal preference.

Before proceeding to our main results, we show that PS^q converges to PS^∞ as $q \rightarrow \infty$. The convergence occurs in all standard metrics; for concreteness, we define the metric by $\|\phi - \hat{\phi}\| := \sup_{\pi \in \Pi, a \in O} |\phi_a(\pi) - \hat{\phi}_a(\pi)|$ for any pair of symmetric random assignments ϕ and $\hat{\phi}$. The convergence of PS^q to PS^∞ is immediate if $\{\Gamma^q\}_{q \in \mathbb{N}}$ are replica economies. In this case, $m_a^q(O') = m_a^\infty(O')$ for all q and a , so the recursive definitions, (3.5), (3.6), (3.7), and (3.8), of the PS procedure for each q -economy all coincide with those of the limiting economy, namely (3.9), (3.10), (3.11), and (3.12). The other cases are established.

Theorem 1. $\|PS^q - PS^\infty\| \rightarrow 0$ as $q \rightarrow \infty$. Further, $PS^q = PS^\infty$ for all $q \in \mathbb{N}$ if $\{\Gamma^q\}_{q \in \mathbb{N}}$ are replica economies.

This theorem shows that PS in the limit economy captures the limiting behavior of PS in a large but finite economy. In this sense, Theorem 1 provides a foundation for a modeling approach that models PS directly in the continuum economy.

4. MAIN RESULT: ASYMPTOTIC EQUIVALENCE

The main purpose of this paper is to investigate how RP and PS compare as the market size grows. Before we present the main result, we offer an example to show that RP and PS can be different even in large finite economies.

Example 3. Consider replica economies $\{\Gamma^q\}_{q \in \mathbb{N}}$ with 2 types of proper object types, a and b , each having quota q in the q -fold replica. Let $N^q = N_\pi^q \cup N_{\pi'}^q$ be the set of agents in the q -fold replica, with N_π^q and $N_{\pi'}^q$ containing $2q$ agents each. Assume that the preferences of the agents are specified by

$$\begin{aligned} (\pi(1), \pi(2), \pi(3)) &= (a, b, \emptyset), \\ (\pi'(1), \pi'(2), \pi'(3)) &= (b, a, \emptyset). \end{aligned}$$

This is the same environment as the one in Example 2. In Examples 1 and 2, we have seen that PS is ordinally efficient in all q -economies, while RP results in an ordinally inefficient random assignment in the base economy.




FIGURE 1. Horizontal axis: Market size q . Vertical axis: $RP_b^q(\pi)$. Note that the assignment coincides with PS (and hence is ordinally efficient) if $RP_b^q(\pi) = 0$.

Figure 1 plots $RP_b^q(\pi)$ as a function of the size of the market q (note that $RP_b^q(\pi) = \|RP^q - PS^q\|$ in this example). $RP_b^q(\pi)$ remains strictly positive even when q is large but finite, so that the ordinal inefficiency of RP does not disappear completely in a finite market. This last observation can be generalized as follows.

Proposition 3. Consider a family $\{\Gamma^q\}_{q \in \mathbb{N}}$ of replica economies. Then, RP^q is ordinally efficient for some $q \in \mathbb{N}$ if and only if $RP^{q'}$ is ordinally efficient for every $q' \in \mathbb{N}$. That is, the random priority assignment is ordinally efficient for all replica economies or ordinally inefficient for all of them.

In particular, Proposition 3 implies that the ordinal inefficiency of RP does not disappear completely in any finitely replicated economy if the random priority assignment is ordinally inefficient in the base economy. On the other hand, in Figure 1 one can see that $\|RP^q - PS^q\|$ decreases as q becomes large in this example. This suggests that the difference between RP and PS may become small as the market size increases and vanishes completely in the limit. The following result shows that this conjecture is true in any sequence of economies (beyond the simple cases of replica economies).

Theorem 2. $\|RP^q - PS^\infty\| \rightarrow 0$ as $q \rightarrow \infty$. Furthermore, $\|RP^q - PS^q\| \rightarrow 0$ as $q \rightarrow \infty$.

We shall give intuition of Theorem 2. The starting point is a recursive formulation of the random priority mechanism given by (3.1) -(3.4). The formulation suggests that the

assignment under the random priority mechanism is similar to the one in the probabilistic serial mechanism, except that in the random priority the random cutoffs replaces the expiration dates in the probabilistic serial mechanism. The basic idea of the proof is to show that the cutoff for each object type in RP converges to the expiration date of that object type in the PS in probability as the size of the market approaches infinity. The convergence will happen if the consumption of each object type in RP during all relevant intervals is close to the corresponding consumption in PS. This will happen under RP in large markets as the law of large number kicks in: with a very high probability, objects are consumed almost proportionately to the number of agents who like that object best among available ones. The proof makes this intuition precise by showing inductively that, with high probability, all cutoffs in RP are sufficiently close to corresponding expiration dates in PS.

5. DISCUSSION AND CONCLUSION

5.1. Random Priority Mechanism in the Limit. Our main result has been established without defining the RP in the limit economy. This omission entails no loss for our purpose, since we are primarily interested in the behavior of a large, but *finite*, economy. Further, defining the RP in the limit economy may require one to describe the aggregate behavior of independent lottery drawing for a continuum of population, and may thus need to appeal to a law of large numbers.¹⁰

There is a way to define the RP in the limit economy, without appealing to a law of large numbers, as has been done by Abdulkadiroğlu, Che, and Yasuda (2008). To do so, we first augment the type of an agent to include his random draw, which is not observed until the random priority is drawn. Formally, a generic agent in the limit economy has type (π, f) representing his preference π and the draw f (which is possibly unobserved by the agent themselves until proper time). The set of agents in the limit economy is represented by the product space $\Pi \times [0, 1]$ endowed with a product measure $\mu \times \nu$, such that $\mu(\pi) = m_\pi^\infty$ for all π and ν is uniform with $\nu([0, f]) = f$ for each $f \in [0, 1]$. In words, the measure of agents with draws less than f is precisely f . This corresponds to the heuristics that the agents in the limit economy obtain random draws in $[0, 1]$ and a law of large number holds for the aggregate distribution (although it is never formally invoked). Again, we assume that a lower draw gives a higher priority for an agent.

¹⁰See Judd (1985) for a classic reference for the associated conceptual problems, and Sun (2006) for a recent treatment.

As with q -economy with $q \in \mathbb{N}$, we characterize the RP^∞ via the cutoff values of the draws for each object. A cutoff \hat{T}_a^∞ for object $a \in O$ is defined such that an agent can obtain a (when he/she wishes) if and only if $f < \hat{T}_a^\infty$. As before, we then define the cutoffs recursively by a sequence of steps. Let $\hat{O}^\infty(0) = \tilde{O}$, $\hat{t}^\infty(0) = 0$, and $\hat{x}_a^\infty(0) = 0$ for every $a \in \tilde{O}$. Given $\hat{O}^\infty(0), \hat{t}^\infty(0), \{\hat{x}_a^\infty(0)\}_{a \in \tilde{O}}, \dots, \hat{O}^\infty(v-1), \hat{t}^\infty(v-1), \{\hat{x}_a^\infty(v-1)\}_{a \in \tilde{O}}$, we let $\hat{t}_\emptyset^\infty := 1$ and for each $a \in O$, define

$$(5.1) \quad \hat{t}_a^\infty(v) = \sup \left\{ t \in [0, 1] \mid \hat{x}_a^\infty(v-1) + m_a^\infty(\hat{O}^\infty(v-1))(t - \hat{t}^\infty(v-1)) < 1 \right\},$$

$$(5.2) \quad \hat{t}^\infty(v) = \min_{a \in \hat{O}^\infty(v-1)} \hat{t}_a^\infty(v),$$

$$(5.3) \quad \hat{O}^\infty(v) = \hat{O}^\infty(v-1) \setminus \{a \in \hat{O}^\infty(v-1) \mid \hat{t}_a^\infty(v) = t^\infty(v)\},$$

$$(5.4) \quad \hat{x}_a^\infty(v) = \hat{x}_a^\infty(v-1) + m_a^\infty(\hat{O}^\infty(v-1))(\hat{t}^\infty(v) - \hat{t}^\infty(v-1)).$$

Comparing (3.9)-(3.12) with (5.1)-(5.4) makes it plainly evident that $\hat{T}_a^\infty = T_a^\infty, \forall a \in O$, with the following conclusion:

Proposition 4. $RP^\infty = PS^\infty$.

This proposition and Theorem 2 imply

Corollary 1. $\|RP^q - RP^\infty\| \rightarrow 0$ as $q \rightarrow \infty$.

Thus RP in the limit economy captures the limiting behavior of RP in a large but finite economy. In this sense, Proposition 4 gives a foundation for a modeling approach that models the random priority mechanism directly in the continuum economy, as has been done, for instance, by Abdulkadiroğlu, Che, and Yasuda (2008).

5.2. Asymptotic Inefficiency in Large Economies. A recurring theme in economics is that large economies can make things “right” in many settings, and our result shares the same theme. Nevertheless, it seems that no single economic insight explains the benefit of large economies. And it is important to investigate what precisely the large economy buys.

To begin, it is often the case the large economy limits individuals’ abilities and incentives to manipulate the system. This is clearly the case in the Walrasian mechanism in exchange economy, as has been shown by Roberts and Postlewaite (1976). It is also the case with the deferred acceptance algorithm in two-sided matching (Kojima and Pathak (2008)) and the probabilistic serial mechanism in one-sided matching (Kojima and Manea (2008)). Even this property is not to be taken for granted, however. The so-called **Boston mechanism** provides an example. The Boston mechanism has been used to place students in public

schools for the K-8 grade. In that mechanism, a school first admits the students who rank it the first, and if, and *only if*, there are seats left, does it admit those who rank it second, and so forth. It is well known that the students have incentives to manipulate the rankings in such a system, and this manipulation incentives do not disappear as the system becomes large.¹¹

Second, one may expect that, with the diminished manipulation incentives, efficiency would be easier to attain in a large economy. The asymptotic ordinal efficiency we find for the RP supports this impression. Yet, even some reasonable mechanisms fail the asymptotic efficiency. Take the case of the **deferred acceptance algorithm with multiple tie-breaking (DA-MTB)**, an adaptation of the celebrated algorithm proposed by Gale and Shapley (1962) to the problem of assigning objects to agents such as student assignment in public schools (see Abdulkadiroğlu, Pathak, and Roth (2005)). In DA-MTB, each object type randomly and independently orders agents and, given the ordering, the assignment is decided by conducting the agent-proposing deferred acceptance algorithm with respect to the submitted preferences and the randomly decided priority profile. It turns out DA-MTB fails even ex post efficiency, let alone ordinal efficiency. Moreover, these inefficiencies do not disappear in the limit economy, as has been shown by Abdulkadiroğlu, Che, and Yasuda (2008).

Third, one plausible conjecture may be that the asymptotic ordinal efficiency is a necessary consequence of a mechanism that produces an ex post efficient assignment in every finite economy. This conjecture turns out to be false. Consider a family $\{\Gamma^q\}$ of replica economies and what we call a **replication-invariant random priority mechanism, $RIRP^q$** , defined as follows. First, in the given q -economy, define a correspondence $\gamma : N^1 \rightarrow N^q$ such that $|\gamma(i)| = q$ for each $i \in N^1$, $\gamma(i) \cap \gamma(j) = \emptyset$ if $i \neq j$, and all agents in $\gamma(i)$ have the same preference as i . Call $\gamma(i)$ i 's clones in the q -fold replica. Let each set $\gamma(i)$ of clones of agent i randomly draws a number f_i independently from a uniform distribution on $[0, 1]$. Second, all the clones with the smallest draw receive their most preferred object, the clones with the second-smallest draw receive their most preferred object from the remaining ones, and so forth. This procedure induces a random assignment. It is clear that $RIRP^q = RP^1$ for any q -fold replica Γ^q . Therefore $\|RIRP^q - RP^1\| \rightarrow 0$ as $q \rightarrow \infty$. Since RP^1 may not be ordinally efficient, the limit random assignment of $RIRP^q$ as $q \rightarrow \infty$ is not ordinally efficient in general.

¹¹See Kojima and Pathak (2008) for a concrete example on this point.

Above all, our analysis shows the equivalence of two different mechanisms beyond showing certain asymptotic properties of given mechanisms. Such an equivalence is not expected even for a large economy, and has few analogues in the literature.

5.3. Existing Priorities. In some applications, the social planner may need to give higher priorities to a subset of agents over others. For example, when allocating graduate dormitory rooms, the housing office at Harvard University assigns rooms to first year students first, and then assign remaining rooms to existing students.

To model such a situation, assume that each student belongs to one of the classes C and, for each $c \in C$, let g_c be a density function over $[0, 1]$. The asymmetric random priority mechanism associated with $g = (g_c)_{c \in C}$ lets each agent i in class c to draw f_i according to the density function g_c independently from others, and the agent with the smallest draw among all agents receives her most preferred object, the agent with the second-smallest draw receives his most preferred object from the remaining ones, and so forth. The random priority mechanism is a special case in which g_c is a uniform distribution on $[0, 1]$ for each $c \in C$.

The asymmetric probabilistic serial mechanism associated with g is defined by simply letting agents in class c to eat with speed $g_c(t)$ at each time $t \in [0, 1]$. The probabilistic serial mechanism is a special case in which g_c is a uniform distribution on $[0, 1]$ for each $c \in C$.

It is not difficult to see that our results generalize to a general profile of distributions g . In particular, given any g , the asymmetric random priority mechanism associated with g and the asymmetric probabilistic serial mechanism associated with g converge to the same limit as $q \rightarrow \infty$.

5.4. Unequal Number of Copies. We focused on a setting in which there are q copies of each object type in the q -economy. It is straightforward to extend our results as long as quotas of object types grows proportionately. More specifically, if there exist positive integers $(q_a)_{a \in O}$ such that the quota of object type a is $q_a q$ in the q -economy, then our result extends with little modification of the proof.

On the other hand, we need *some* assumption about the growth rate of quotas. Suppose that, for instance, quotas of some objects are q but quotas of others stay at one. Then, one can easily create an example in which random priority assignment of objects with quota one does not converge to those under the probabilistic serial mechanism. However, such an example does not seem to be a large problem, since in the large market, assignment of object types with small quotas has only limited influence on overall welfare in the economy.

5.5. Multiple Unit Assignments. Consider a generalization of our basic setting, in which each agent can obtain multiple units of objects. More specifically, we assume that there is a fixed integer k such that each agent can receive k objects. When $k = 1$, the model reduces to the model of the current paper. Assignment of popular courses in schools is one example of such a multiple unit assignment problem. See, for example, Kojima (2008) for formal definition of the model.

We consider two generalizations of the random priority mechanism to the current setting. In the **once-and-for-all random priority mechanism**, each agent i randomly draws a number f_i independently from a uniform distribution on $[0, 1]$ and, given the ordering, the agent with the smallest draw receives her most preferred k objects, the agent with the second-smallest draw receives his most preferred k objects from the remaining ones, and so forth. In the **draft random priority mechanism**, each agent i randomly draws a number f_i independently from a uniform distribution on $[0, 1]$. Second, the agent with the smallest draw receives her most preferred object, the agent with the second-smallest draw receives his most preferred object from the remaining ones, and so forth. Then agents obtain a random draw again and repeat the procedure k times.

We introduce two generalizations of the probabilistic serial mechanism. In the **multiunit-eating probabilistic serial** mechanism, each agent “eats” her k most preferred available objects with speed one at every time $t \in [0, 1]$. In the **one-at-a-time probabilistic serial** mechanism, each agent “eats” the best available object with speed one at every time $t \in [0, k]$.

Our analysis can be adapted to this situation to show that the once-and-for-all random priority mechanism converges to the same limit as the multiunit-eating probabilistic serial mechanism, whereas the draft random priority mechanism converges to the same limit as the one-at-a-time probabilistic serial mechanism.

It is easy to see that the multiunit-eating probabilistic serial mechanism may not be ordinally efficient, while the one-at-a-time probabilistic serial mechanism is ordinally efficient. This may give an insight to some issues in multiple unit assignment. It is well known that the once-and-for-all random priority mechanism is ex post efficient, but the mechanism is rarely used in practice. Rather, the draft mechanism is often used in application, for instance sports drafting and allocations of courses in business schools. One of the reasons may be that the once-and-for-all random priority mechanism is ordinally inefficient even in the limit economy, whereas the draft random priority mechanism converges to an ordinally efficient mechanism as the economy becomes large, as in course allocation in schools.

5.6. Concluding Remarks. The random priority (random serial dictatorship) mechanism is a common method to assign objects to individuals, while it is recently challenged by a new probabilistic serial mechanism. We showed that these mechanisms become equivalent when the market becomes large. More specifically, given a set of object types, the random assignments in these mechanisms converge to each other as the number of copies of each object type approaches infinity. Thus inefficiency of the random priority mechanism becomes small in large markets.

Our equivalence is asymptotic and the random priority and the probabilistic serial mechanisms do not exactly coincide in large but finite economies. It would be interesting to see how these competing mechanisms perform in assignment problems in real life such as student placement in public schools.

APPENDIX

A. PROOF OF THEOREM 1

It suffices to show that $\sup_{a \in O} |T_a^q - T_a^\infty| \rightarrow 0$ as $q \rightarrow \infty$. To this end, let

$$(A1) \quad L > \max \left\{ 5, 2 \max \left\{ \frac{1}{m_a^\infty(O')} \vee m_a^\infty(O') \mid O' \subset O, a \in O', m_a^\infty(O') > 0 \right\} \right\},$$

and let $K := \min\{1 - x_a^\infty(v) \mid a \in O^\infty(v), v < \bar{v}^\infty\} > 0$.

Fix any $\epsilon > 0$ such that

$$(A2) \quad 2L^{4\bar{v}^\infty} \epsilon < \min\left\{ \min_{v \in 1, \dots, \bar{v}^\infty} |t^\infty(v) - t^\infty(v-1)|, K \right\}.$$

By assumption there exists Q such that, for each $q > Q$,

$$(A3) \quad |m_a^q(O') - m_a^\infty(O')| < \epsilon, \forall O' \subset \tilde{O}, \forall a \in O'.$$

Fix any such q . We show that $T_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$. To this end, we argue recursively. Suppose for any $v' \leq v-1$, $T_a^q \in (t^\infty(v') - L^{4v'}\epsilon, t^\infty(v') + L^{4v'}\epsilon)$ if and only if $t_a^\infty(v') = t^\infty(v')$, and further that, for each $a \in O^\infty(v-1)$, $x_a^q(k) \in (x_a^\infty(v-1) - L^{4(v-1)}\epsilon, x_a^\infty(v-1) + L^{4(v-1)}\epsilon)$, where k is the largest element of $J_{v-1} := \{i \mid \text{there exists } a \text{ s.t. } t_a^q(i) = t^q(i) \text{ and } T_a^\infty = t^\infty(v-1)\}$. We shall then prove that $T_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$, and that, for each $a \in O^\infty(v)$, $x_a^q(l) \in (x_a^\infty(v) - L^{4v}\epsilon, x_a^\infty(v) + L^{4v}\epsilon)$, where l is the largest element of J_v .

Let k be the largest element of J_{v-1} . It then follows that $O^q(k) = O^\infty(v-1)$.

Claim 1. For any $a \in O^\infty(v-1)$, $t_a^q(k+1) > t^\infty(v) - L^{4v-2}\epsilon$.

Proof. Let a be the first good to expire in $O^\infty(v-1)$ under PS^q . Assume, for contradiction, that

$$(A4) \quad t_a^q(k+1) \leq t^\infty(v) - L^{4v-2}\epsilon.$$

Recall, by inductive assumption, that

$$(A5) \quad x_a^q(k) < x_a^\infty(v-1) + L^{4(v-1)}\epsilon.$$

Thus,

$$\begin{aligned} x_a^q(k+1) &= x_a^q(k) + m_a^q(O^q(k))(t_a^q(k+1) - t_a^q(k)) \\ &\leq x_a^q(k) + m_a^q(O^q(k))(t^\infty(v) - L^{4v-2}\epsilon - t^\infty(v-1) + L^{4(v-1)}\epsilon) \\ &\leq x_a^q(k) + m_a^q(O^q(k))[t^\infty(v) - t^\infty(v-1) - L^{4v-3}\epsilon] \\ (A6) \quad &< x_a^\infty(v-1) + L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) - L^{4v-3}\epsilon] + \epsilon, \end{aligned}$$

where the first equality follows from definition of PS^q (3.8), the first inequality follows from the inductive assumption and (A4), the second inequality follows from (A1), and the third inequality follows from (A2), (A3) and (A5).

There are two cases. Suppose first $m_a^\infty(O^\infty(v-1)) = 0$. Then, the last line of (A6) becomes

$$x_a^\infty(v-1) + L^{4(v-1)}\epsilon + \epsilon,$$

which is strictly less than 1, since $a \in O^\infty(v-1)$ and since (A2) holds. Suppose next $m_a^\infty(O^\infty(v-1)) > 0$. Then, the last line of (A6) equals

$$\begin{aligned} &x_a^\infty(v-1) + L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) - L^{4v-3}\epsilon] + \epsilon \\ &< x_a^\infty(v-1) + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1)] \\ &\leq 1, \end{aligned}$$

where the first inequality follows from (A1), and the second follows since $a \in O^\infty(v-1)$. In either case, we have a contradiction to the fact that a expires at step $k+1$. \parallel

Claim 2. Let b be the object that expires the last in the q -economy among the set $\{a \in O^\infty(v-1) | t_a^\infty(v) = t^\infty(v)\}$. If b expires at stage $l \geq k+1$ in the q -economy, then $t_b^q(l) \leq t^\infty(v) + L^{4v-2}\epsilon$.

Proof. The result follows since

$$\begin{aligned}
t_b^q(l) &\leq (1 - x_b^q(l-1)) \frac{1}{m_b^q(O^q(l-1))} + t^q(l-1) \\
&= (1 - x_b^q(l-1)) \frac{1}{m_b^q(O^q(l-1))} + \left(\sum_{k+1 \leq j \leq l-1} (t^q(j) - t^q(j-1)) \right) + t^q(k) \\
&= (1 - x_b^q(l-1)) \frac{1}{m_b^q(O^q(l-1))} + \left(\sum_{k+1 \leq j \leq l-1} (x_b^q(j) - x_b^q(j-1)) \frac{1}{m_b^q(O^q(j-1))} \right) + t^q(k) \\
&\leq (1 - x_b^q(k)) \frac{1}{m_b^q(O^q(k))} + t^q(k) \\
&< t_b^\infty(v) + L^{4v-2}\epsilon \\
&= t^\infty(v) + L^{4v-2}\epsilon.
\end{aligned}$$

The first inequality follows from (3.5), the second equality follows from (3.8), and the second inequality holds since $m_b^q(O^q(j)) \geq m_b^q(O^q(i))$ if $j > i$ and $b \in O^q(j)$; the third inequality follows from an argument analogous to the ones used above; and the last equality follows since $t_b^\infty(v) = t^\infty(v)$. \parallel

Claim 3. If $a \in O^\infty(v)$ and $v < \bar{v}^\infty$, then $T_a^q > t^\infty(v) + L^{4v}\epsilon$.

Proof. Suppose otherwise. Let c be the object in $O^\infty(v)$ that expires the first in the q -economy. Let j be the step at which it expires. Then, we must have $t_c^q(j) < 1$ and $x_c^q(j) = 1$. Since c is the first object to expire in $O^\infty(v)$, at each of steps $k+1, \dots, j-1$, some good in $O^\infty(v-1) \setminus O^\infty(v) = \{a \in O^\infty(v-1) \mid t_a^\infty(v) = t^\infty(v)\}$ expires. (If $j = k+1$, then no other object expires in between step k and step j .) By the previous analysis, this implies $t^q(k+1) > t^\infty(v) - L^{4v-2}\epsilon$. Therefore,

$$\begin{aligned}
x_c^q(j) &= x_c^q(k) + \sum_{i=k+1}^j m_c^q(O^q(i-1))(t^q(i) - t^q(i-1)) \\
&\leq x_c^q(k) + m_c^q(O^q(k))(t^q(k+1) - t^q(k)) + m_c^q(O^q(j-1))(t^q(j) - t^q(k+1)) \\
&\leq x_c^\infty(v-1) + L^{4(v-1)}\epsilon + (m_c^\infty(O^q(k)) + \epsilon) ((t^\infty(v) + L^{4v-2}\epsilon) - (t^\infty(v-1) - L^{4(v-1)}\epsilon)) \\
&\quad + (m_a^\infty(O^q(j)) + \epsilon) (L^{4v}\epsilon - L^{4v-2}\epsilon) \\
&\leq x_c^\infty(v) + L^{4v+1}\epsilon \\
&\leq 1 - K + L^{4\bar{v}^\infty}\epsilon \\
&< 1,
\end{aligned}$$

which contradicts the assumption that c expires at step j . \parallel

The arguments so far prove that $T_a^q \in (t^\infty(v) - L^{4v-2}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \subset (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $T_a^\infty = t^\infty(v)$. It now remains to prove the following:

Claim 4. For each $a \in O^\infty(v)$, $x_a^q(l) \in (x_a^\infty(v) - L^{4v}\epsilon, x_a^\infty(v) + L^{4v}\epsilon)$, where l is the largest element of J_v .

Proof. Fix any $a \in O^\infty(v)$. Then,

$$\begin{aligned}
x_a^q(l) &= x_a^q(k) + \sum_{j=k+1}^l m_a^q(O^q(j-1))(t^q(j) - t^q(j-1)) \\
&\leq x_a^q(k) + m_a^q(O^q(k))(t^q(k+1) - t^q(k)) + m_a^q(O^q(l))(t^q(l) - t^q(k+1)) \\
&\leq x_a^\infty(v-1) + L^{4(v-1)}\epsilon + (m_a^\infty(O^q(k)) + \epsilon)(t^\infty(v) - t^\infty(v-1) + 2L^{4v-2}\epsilon) \\
&\quad + (m_a^\infty(O^q(l)) + \epsilon)(2L^{4v-2}\epsilon) \\
&< x_a^\infty(v-1) + (m_a^\infty(O^\infty(v-1)))(t^\infty(v) - t^\infty(v-1)) + L^{4v}\epsilon \\
&= x_a^\infty(v) + L^{4v}\epsilon.
\end{aligned}$$

A symmetric argument yields $x_a^q(l) \geq x_a^\infty(v) - L^{4v}\epsilon$. \parallel

We have thus completed the recursive argument, which taken together proves that $T_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$, for any $q > Q$ for some $Q \in \mathbb{N}$. Since $\epsilon > 0$ can be arbitrarily small, $T_a^q \rightarrow T_a^\infty$ as $q \rightarrow \infty$. Since there are only a finite number of objects and a finite number of preference types, $\|PS^q - PS^\infty\| \rightarrow 0$ as $q \rightarrow \infty$.

B. PROOF OF THEOREM 2

Let L be a real number satisfying

$$(B1) \quad L > \max\{5, 2 \max\{\frac{1}{m_a^\infty(O')} \vee m_a^\infty(O') \mid O' \subset O, a \in O', m_a^\infty(O') > 0\}\},$$

and let $K := \min\{1 - x_a^\infty(v) \mid a \in O^\infty(v), v < \bar{v}^\infty\} > 0$.

Fix an agent i_0 of preference type $\pi_0 \in \Pi$ and consider a random assignment for agents of type π_0 . Consider the following events:¹²

$$\begin{aligned}
E_1^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v-1) - L^{4(v-1)}\epsilon, t_a^\infty(v) - L^{4v-2}\epsilon) < & m_\pi^\infty[t_a^\infty(v) - t_a^\infty(v-1) - L^{4v-3}\epsilon], \\
E_2^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v-1) + L^{4(v-1)}\epsilon, t_a^\infty(v) + L^{4v-2}\epsilon) \geq & m_\pi^\infty[t_a^\infty(v) - t_a^\infty(v-1) + L^{4v-3}\epsilon], \quad v \neq \bar{v}^\infty, \\
E_2^q(\bar{v}^\infty, \pi) &: & \{f_{-i_0} \in [0, 1]^{|N^q-1|}\}, \\
E_3^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v-1) - L^{4(v-1)}\epsilon, t_a^\infty(v) + L^{4v-2}\epsilon) < & m_\pi^\infty[t_a^\infty(v) - t_a^\infty(v-1) + 2L^{4v-2}\epsilon], \\
E_4^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v) - L^{4v-2}\epsilon, t_a^\infty(v) + L^{4v}\epsilon) < & m_\pi^\infty \times 2L^{4v}\epsilon, \\
E_5^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v) - L^{4v-2}\epsilon, t_a^\infty(v) + L^{4v-2}\epsilon) < & m_\pi^\infty \times 3L^{4v-2}\epsilon, \\
E_6^q(v, \pi) &: \hat{m}_\pi^q(t_a^\infty(v-1) + L^{4(v-1)}\epsilon, t_a^\infty(v) - L^{4v-2}\epsilon) \geq & m_\pi^\infty[t_a^\infty(v) - t_a^\infty(v-1) - L^{4v-2}\epsilon].
\end{aligned}$$

Lemma 1. For any $\epsilon > 0$ such that

$$(B2) \quad 2L^{4\bar{v}^\infty}\epsilon < \min\left\{\min_{v \in \{1, \dots, \bar{v}^\infty\}}\{t^\infty(v) - t^\infty(v-1)\}, K\right\},$$

there exists Q such that the following is true for any $q > Q$: If the realization of f_{-i_0} is such that events $E_1^q(v, \pi)$, $E_2^q(v, \pi)$, $E_3^q(v, \pi)$, $E_4^q(v, \pi)$, $E_5^q(v, \pi)$ and $E_6^q(v, \pi)$ hold for all $v \in \{1, \dots, \bar{v}^\infty\}$ and $\pi \in \Pi$ with $m_\pi^\infty > 0$, then $\hat{T}_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$.

Proof. There exists Q such that

$$(B3) \quad \sum_{\pi \in \Pi: m_\pi^\infty = 0} m_\pi^q < \epsilon,$$

for any $q > Q$. Fix any such q and suppose that the realization of f_{-i_0} is such that $E_1^q(v, \pi)$, $E_2^q(v, \pi)$, $E_3^q(v, \pi)$, $E_4^q(v, \pi)$, $E_5^q(v, \pi)$ and $E_6^q(v, \pi)$ hold for all v, π with $m_\pi^\infty > 0$ as described in the statement of the Lemma. We show the lemma inductively. Suppose for any $v' \leq v-1$, $\hat{T}_a^q \in (t^\infty(v') - L^{4v'}\epsilon, t^\infty(v') + L^{4v'}\epsilon)$ if and only if $t_a^\infty(v') = t^\infty(v')$, and further that, for each $a \in O^\infty(v-1)$, $\hat{x}_a^q(k) \in (x_a^\infty(v-1) - L^{4(v-1)}\epsilon, x_a^\infty(v-1) + L^{4(v-1)}\epsilon)$, where k is the largest element of $\hat{J}_{v-1} := \{i \mid \text{there exists } a \text{ s.t. } \hat{t}_a^q(i) = \hat{t}^q(i) \text{ and } T_a^\infty = t^\infty(v-1)\}$. We shall then prove that $\hat{T}_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$, and that, for each $a \in O^\infty(v)$, $\hat{x}_a^q(l) \in (x_a^\infty(v) - L^{4v}\epsilon, x_a^\infty(v) + L^{4v}\epsilon)$, where l is the largest element of \hat{J}_v .

Let k be the largest element of \hat{J}_{v-1} . It then follows that $\hat{O}^q(k) = O^\infty(v-1)$. Fix any $a \in O^\infty(v-1)$.

¹² $E_2^q(\bar{v}^\infty, \pi)$ holds for any realization of ordering. The notation is introduced only for expositional simplicity in what follows.

Claim 5. $\hat{t}_a^q(k+1) > t^\infty(v) - L^{4v-2}\epsilon$ for all $a \in O^\infty(v-1)$.

Proof. Let a be the first good to expire in $O^\infty(v-1)$ under RP^q . Assume, for contradiction, that

$$(B4) \quad \hat{t}_a^q(k+1) \leq t^\infty(v) - L^{4v-2}\epsilon.$$

Recall, by inductive assumption, that

$$(B5) \quad \hat{x}_a^q(k) < x_a^\infty(v-1) + L^{4(v-1)}\epsilon.$$

Thus,

$$(B6) \quad \begin{aligned} \hat{x}_a^q(k+1) &= \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); \hat{t}^q(k), \hat{t}_a^q(k+1)) \\ &\leq \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); t^\infty(v-1) - L^{4(v-1)}\epsilon, t^\infty(v) - L^{4v-2}\epsilon) \\ &< x_a^\infty(v-1) + L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) - L^{4v-3}\epsilon] + \epsilon, \end{aligned}$$

where the first equality follows from (3.4) in the definition of RP^q , the first inequality follows from the inductive assumption and (B4), and the second inequality follows from the assumption that $E_1^q(v, \pi)$ holds and conditions (B3) and (B5).

There are two cases. Suppose first $m_a^\infty(O^\infty(v-1)) = 0$. Then, the last line of (B6) becomes

$$x_a^\infty(v-1) + L^{4(v-1)}\epsilon + \epsilon,$$

which is strictly less than 1, since $a \in O^\infty(v-1)$ and since (B2) holds. Suppose next $m_a^\infty(O^\infty(v-1)) > 0$. Then, the last line of (B6) equals

$$\begin{aligned} &x_a^\infty(v-1) + L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) - L^{4v-3}\epsilon] + \epsilon \\ &< x_a^\infty(v-1) + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1)] \\ &\leq 1, \end{aligned}$$

where the first inequality follows from (B1), and the second follows since $a \in O^\infty(v-1)$. In either case, we have a contradiction to the fact that a expires at step $k+1$. \parallel

Claim 6. Let a be the object that expires the last, in RP^q with the given order f , in the q -economy among the set $\{b \in O^\infty(v-1) | t_b^\infty(v) = t^\infty(v)\}$. Suppose a expires at stage $l \geq k+1$ in the q -economy. Then,

$$(B7) \quad \hat{t}^q(l) < t^\infty(v) + L^{4v-2}\epsilon.$$

Proof. Suppose, for contradiction, that

$$(B8) \quad \hat{t}^q(l) \geq t^\infty(v) + L^{4v-2}\epsilon.$$

Then,

$$\begin{aligned} \hat{x}_a^q(l) &= \hat{x}_a^q(k) + \sum_{j=k+1}^l \hat{m}_a^q(\hat{O}^q(j-1); \hat{t}^q(j-1), \hat{t}^q(j)) \\ &\geq \hat{x}_a^q(k) + \sum_{j=k+1}^l \hat{m}_a^q(\hat{O}^q(k); \hat{t}^q(j-1), \hat{t}^q(j)) \\ &= \hat{x}_a^q(k) + \hat{m}_a^q(O^\infty(v-1); \hat{t}^q(k), \hat{t}^q(l)) \\ &> x_a^\infty(v-1) - L^{4(v-1)}\epsilon + \hat{m}_a^q(O^\infty(v-1); t^\infty(v-1) + L^{4(v-1)}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \\ &\geq x_a^\infty(v-1) - L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) + L^{4v-3}\epsilon] \\ &> x_a^\infty(v-1) + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1)] \\ &= x_a^\infty(v) = 1, \end{aligned}$$

where the first equality follows from (3.4), the first inequality follows since $\hat{m}_a^q(\hat{O}^q(j-1); t, t') \geq m_a^q(\hat{O}^q(k); t, t')$ for any $j \geq k+1$ and $t \leq t'$ by $\hat{O}^q(j-1) \subseteq \hat{O}^q(k)$, the second equality follows from $\hat{O}^q(k) = O^\infty(v-1)$ and the definition of \hat{m}_a^q , the second inequality follows from the inductive assumption and (B8), the third inequality follows from the assumption that $E_2^q(v, \pi)$ holds, and the fourth inequality follows from (B1). \parallel

Claim 7. If $a \in O^\infty(v)$ and $v < \bar{v}^\infty$, then $\hat{T}_a^q > t^\infty(v) + L^{4v}\epsilon$.

Proof. Suppose otherwise. Let c be the object in $O^\infty(v)$ that expires the first in the q -economy. Let j be the step at which it expires. Then, we must have

$$(B9) \quad \hat{t}_c^q(j) \leq t^\infty(v) + L^{4v}\epsilon,$$

and $\hat{x}_c^q(j) = 1$. Since c is the first object to expire in $O^\infty(v)$, at each of steps $k+1, \dots, j-1$, some good in $O^\infty(v-1) \setminus O^\infty(v) = \{a \in O^\infty(v-1) \mid t_a^\infty(v) = t^\infty(v)\}$ expires. (If $j = k+1$, then no other object expires in between step k and step j .) By Claim 5, this implies

$\hat{t}^q(k+1) > t^\infty(v) - L^{4v-2}\epsilon$. Therefore,

$$\begin{aligned}
\hat{x}_c^q(j) &= \hat{x}_c^q(k) + \sum_{i=k+1}^j \hat{m}_c^q(\hat{O}^q(i-1); \hat{t}^q(i-1), \hat{t}^q(i)) \\
&\leq \hat{x}_c^q(k) + \hat{m}_c^q(\hat{O}^q(k); \hat{t}^q(k), \hat{t}^q(k+1)) + \hat{m}_c^q(\hat{O}^q(j-1); \hat{t}^q(k+1), \hat{t}^q(j)) \\
&\leq \hat{x}_c^q(k) + \hat{m}_c^q(\hat{O}^q(k); t^\infty(v-1) - L^{4(v-1)}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \\
&\quad + \hat{m}_c^q(\hat{O}^q(j-1); t^\infty(v) - L^{4v-2}\epsilon, t^\infty(v) + L^{4v}\epsilon) \\
&\leq x_c^\infty(v-1) + L^{4(v-1)}\epsilon + m_c^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) + 2L^{4v-2}\epsilon] \\
&\quad + m_c^\infty(\hat{O}^q(j-1) \times 2L^{4v}\epsilon + \epsilon) \\
&\leq x_c^\infty(v) + L^{4v+1}\epsilon \\
&\leq 1 - K + L^{4\bar{v}^\infty}\epsilon \\
&< 1,
\end{aligned}$$

where the first equality follows from (3.4), the first inequality follows since $\hat{m}_c^q(\hat{O}^q(j-1); t, t') \geq m_c^q(\hat{O}^q(i-1); t, t')$ for any $j \geq i$ by $\hat{O}^q(j-1) \subseteq \hat{O}^q(i-1)$, the second inequality follows from the inductive assumption, Claims 5 and 6, the third inequality follows from the inductive assumption, $E_3^q(v, \pi)$, $E_4^q(v, \pi)$ and (B3), the fourth inequality follows from (3.12) and (B1), the fifth inequality follows from the definition of K , and the last inequality follows from the assumption that $2L^{4\bar{v}^\infty}\epsilon < K$. Thus we obtain $\hat{x}_c^q(j) < 1$, which contradicts the assumption that c expires at step j . \parallel

The arguments so far prove that $\hat{T}_a^q \in (t^\infty(v) - L^{4v-2}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \subset (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $T_a^\infty = t^\infty(v)$. It now remains to show the following.

Claim 8. For each $a \in O^\infty(v)$, $x_a^q(l) \in (x_a^\infty(v) - L^{4v}\epsilon, x_a^\infty(v) + L^{4v}\epsilon)$, where l is the largest element of \hat{J}_v .

Proof. Fix any $a \in O^\infty(v)$. Then,

$$\begin{aligned}
\hat{x}_a^q(l) &= \hat{x}_a^q(k) + \sum_{j=k+1}^l \hat{m}_a^q(\hat{O}^q(j-1); \hat{t}^q(j-1), \hat{t}^q(j)) \\
&\leq \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); \hat{t}^q(k), \hat{t}^q(k+1)) + \hat{m}_a^q(\hat{O}^q(l); \hat{t}^q(k+1), \hat{t}^q(l)) \\
&\leq \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); t^\infty(v-1) - L^{4(v-1)}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \\
&\quad + \hat{m}_a^q(\hat{O}^q(l); t^\infty(v) - L^{4v-2}\epsilon, t^\infty(v) + L^{4v-2}\epsilon) \\
&< x_a^\infty(v-1) + L^{4(v-1)}\epsilon + m_a^\infty(\hat{O}^q(k))(t^\infty(v) - t^\infty(v-1) + 2L^{4v-2}\epsilon) \\
&\quad + m_a^\infty(\hat{O}^q(l)) \times 3L^{4v-2}\epsilon + 2\epsilon \\
&< x_a^\infty(v-1) + (m_a^\infty(O^\infty(v-1)))(t^\infty(v) - t^\infty(v-1)) + L^{4v}\epsilon \\
&= x_a^\infty(v) + L^{4v}\epsilon,
\end{aligned}$$

where the first equality follows from (3.4), the first inequality follows from $m_a^q(\hat{O}^q(l); t, t') \geq m_a^q(\hat{O}^q(j); t, t')$ for all $l \geq j$, the second inequality follows from the inductive assumption and Claims 5 and 6, the third inequality follows from the inductive assumption, (B3) and $E_3^q(v, \pi)$ and $E_5^q(v, \pi)$, the fourth inequality follows from $\hat{O}^q(k) = O^\infty(v-1)$ and (B1), and the last inequality follows from (3.12).

Next we obtain

$$\begin{aligned}
\hat{x}_a^q(l) &= \hat{x}_a^q(k) + \sum_{j=k+1}^l \hat{m}_a^q(\hat{O}^q(j-1); \hat{t}^q(j-1), \hat{t}^q(j)) \\
&\geq \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); \hat{t}^q(k), \hat{t}^q(l)) \\
&\geq \hat{x}_a^q(k) + \hat{m}_a^q(\hat{O}^q(k); t^\infty(v-1) + L^{4(v-1)}\epsilon, t^\infty(v) - L^{4v-2}\epsilon) \\
&\geq x_a^\infty(v-1) - L^{4(v-1)}\epsilon + m_a^\infty(O^\infty(v-1))[t^\infty(v) - t^\infty(v-1) - 2L^{4v-2}\epsilon] \\
&> x_a^\infty(v) - L^{4v}\epsilon,
\end{aligned}$$

where the first inequality follows from $\hat{O}^q(j-1) \subseteq \hat{O}^q(k)$ for any $j \geq k+1$, the second inequality follows from the inductive assumption and Claim 5, the third inequality follows from the inductive assumption and $E_6^q(v, \pi)$, and the last inequality follows from (3.12) and (B1). These inequalities complete the proof. \parallel

We have thus completed the recursive argument, which taken together proves that $\hat{T}_a^q \in (t^\infty(v) - L^{4v}\epsilon, t^\infty(v) + L^{4v}\epsilon)$ if and only if $t_a^\infty(v) = t^\infty(v)$, for any $q > Q$ for some $Q \in \mathbb{N}$. \square

Proof of Theorem 2. We shall show that for any $\varepsilon > 0$ there exists Q such that, for any $q > Q$, for any $\pi_0 \in \Pi$ and $a \in O$,

$$(B10) \quad |PS_a^\infty(\pi_0) - RP_a^q(\pi_0)| < (2L^{4(n+1)} + 6n(n+1)(n+1)!) \varepsilon.$$

Since n is a finite constant, relation (B10) implies the Theorem.

To show this first assume, without loss of generality, that ε satisfies (B2) and Q is so large that (B3) holds for any $q > Q$. We have

$$\begin{aligned}
RP_a^q(\pi_0) &= E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \right] \\
&= E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \times Pr \left[\bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \\
&+ E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \overline{\bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi)} \right] \times Pr \left[\overline{\bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi)} \right] \\
&= E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \times \left(1 - Pr \left[\bigcup_{i=1}^6 \bigcup_{v=1}^{\bar{v}^\infty} \bigcup_{\pi \in \Pi: m_\pi^\infty > 0} \overline{E_i^q(v, \pi)} \right] \right) \\
&+ E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \overline{\bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi)} \right] \times Pr \left[\bigcup_{i=1}^6 \bigcup_{v=1}^{\bar{v}^\infty} \bigcup_{\pi \in \Pi: m_\pi^\infty > 0} \overline{E_i^q(v, \pi)} \right] \\
&= E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \\
&+ \left\{ E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \overline{\bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi)} \right] - E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \right\} \\
(B11) \quad &\times Pr \left[\bigcup_{i=1}^6 \bigcup_{v=1}^{\bar{v}^\infty} \bigcup_{\pi \in \Pi: m_\pi^\infty > 0} \overline{E_i^q(v, \pi)} \right],
\end{aligned}$$

where for any event E , \bar{E} is the complement event of E .

First we bound the first term of expression (B11). Since $\bar{v}^\infty \leq n+1$, Lemma 1 implies that

$$E \left[\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \middle| \bigcap_{i=1}^6 \bigcap_{v=1}^{\bar{v}^\infty} \bigcap_{\pi \in \Pi: m_\pi^\infty > 0} E_i^q(v, \pi) \right] \in [T_a^\infty - \tau_a^\infty(\pi_0) - 2L^{4(n+1)}\varepsilon, T_a^\infty - \tau_a^\infty(\pi_0) + 2L^{4(n+1)}\varepsilon].$$

Second, we bound the second term of expression (B11). By the weak law of large numbers, for any $\varepsilon > 0$, there exists Q such that $Pr \left[\overline{E_i^q(v, \pi)} \right] < \varepsilon$ for any $i \in \{1, 2, 3, 4, 5, 6\}$, $q > Q$, $v \in \{1, \dots, \bar{v}^\infty\}$ and $\pi \in \Pi$ with $m_\pi^\infty > 0$. Since there are at most $6(n+1)(n+1)!$ such events and, in general, the sum of probabilities of a number of events is weakly larger than the probability of the union of the events (Boole's inequality), we obtain

$$\begin{aligned} Pr \left[\bigcup_{i=1}^6 \bigcup_{v=1}^{\bar{v}^\infty} \bigcup_{\pi \in \Pi: m_\pi^\infty > 0} \overline{E_i^q(v, \pi)} \right] &\leq \sum_{i=1}^6 \sum_{v=1}^{\bar{v}^\infty} \sum_{\pi \in \Pi: m_\pi^\infty > 0} Pr \left[\overline{E_i^q(v, \pi)} \right] \\ &\leq 6(n+1)(n+1)!\varepsilon. \end{aligned}$$

Since $\hat{T}_a^q - \hat{\tau}_a^q(\pi_0) \in [0, 1]$ for any a, q and π_0 , the second term of equation (B11) is in $[-6(n+1)(n+1)!\varepsilon, 6(n+1)(n+1)!\varepsilon]$.

From the above arguments, we have that

$$|PS_a^\infty(\pi_0) - RP_a^q(\pi_0)| < (2L^{4(n+1)} + 6(n+1)(n+1)!\varepsilon),$$

completing the proof. \square

C. PROOF OF PROPOSITION 3

The proposition uses the following two lemmas. Let $\{\Gamma^q\}$ be a family of replica economies. Given any q , define a correspondence $\gamma : N^1 \rightarrow N^q$ such that $|\gamma(i)| = q$ for each $i \in N^1$, $\gamma(i) \cap \gamma(j) = \emptyset$ if $i \neq j$, and all agents in $\gamma(i)$ have the same preference as i . Call $\gamma(i)$ i 's clones in the q -fold replica.

Lemma 2. For all $q \in \mathbb{N}$ and $a, b \in \tilde{O}$, $a \triangleright (RP^1, m^1) b \iff a \triangleright (RP^q, m^q) b$.

Proof. We proceed in two steps.

(i) $a \triangleright (RP^1, m^1) b \implies a \triangleright (RP^q, m^q) b$: Suppose first $a \triangleright (RP^1, m^1) b$. There exists an individual $i^* \in N^1$ and an ordering $(i_{(1)}^1, \dots, i_{(|N^1|)}^1)$ (implied by some draw $f^1 \in [0, 1]^{|N^1|}$) such that the agents in front of i^* in that ordering consume all the goods that i^* prefers to b but not b , and i^* consumes b .

Now consider the q -fold replica. With positive probability, we have an ordering $(\bar{\gamma}(i_{(1)}^1), \dots, \bar{\gamma}(i_{(|N^1|)}^1))$, where $\bar{\gamma}(i)$ is an arbitrary permutation of $\gamma(i)$. Under this ordering, each agent in $\gamma(i_{(j)}^1)$ will consume a copy of the good agent $i_{(j)}^1$ in the base economy will consume, and all the agents in $\gamma(i^*)$ will consume b (despite preferring a over b). This proves that $a \triangleright (RP^q, m^q) b$.

(ii) $a \triangleright (RP^q, m^q) b \implies a \triangleright (RP^1, m^1) b$: Suppose $a \triangleright (RP^q, m^q) b$. Then, with positive probability, a draw $f^q \in [0, 1]^{|N^q|}$ entails an ordering in which the agents ahead of $i^* \in N^q$ consume all of the goods that i^* prefers to b , but not all the copies of b have been consumed

by them. List these goods in the order that their last copies are consumed, and let the set of these goods be $\hat{O} := \{o_1, \dots, o_m\} \subset O$, where o_l is completely consumed before o_{l+1} for all $l = 1, \dots, m-1$. (Note that $a \in \hat{O}$.) Let i^{**} be such that $i^* \in \gamma(i^{**})$.

We first construct a correspondence $\xi : \hat{O} \mapsto N^1 \setminus \{i^{**}\}$ defined by

$$\xi(o) := \{i \in N^1 \setminus \{i^{**}\} \mid \exists j \in \gamma(i) \text{ who consumes } o \text{ under } f^q\}.$$

Claim 9. Any agent in N^q who consumes o_l prefers o_l to all goods in $\tilde{O} \setminus \{o_1, \dots, o_{l-1}\}$ under f^q . Hence, any agent in $\xi(o_l)$ prefers o_l to all goods in $\tilde{O} \setminus \{o_1, \dots, o_{l-1}\}$.

Claim 10. For each $O' \subset \hat{O}$, $|\cup_{o \in O'} \xi(o)| \geq |O'|$.

Proof. Suppose otherwise. Then, there exists $O' \subset \hat{O}$ such that $k := |\cup_{o \in O'} \xi(o)| < |O'| =: l$. Reindex the sets so that $\cup_{o \in O'} \xi(o) = \{a^1, \dots, a^k\}$ and $O' = \{o^1, \dots, o^l\}$. Let x_{ij} denote the number of clones of agent $a^j \in \xi(o^i)$ who consume o^i in the q -fold replica under f^q .

Since $\sum_{i=1}^l x_{ij} \leq |\gamma(a^j)| = q$,

$$\sum_{j=1}^k \sum_{i=1}^l x_{ij} \leq kq.$$

At the same time, all q copies of each good in O' are consumed, and at most $q-1$ clones of i^{**} could be those contributing to that consumption. Therefore,

$$\sum_{i=1}^l \sum_{j=1}^k x_{ij} \geq lq - (q-1) = (l-1)q + 1 > kq,$$

We thus have a contradiction. \parallel

By Hall's Theorem, Claim 10 implies that there exists a mapping $\mu : \hat{O} \mapsto N^1 \setminus \{i^{**}\}$ such that $\mu(o) \in \xi(o)$ for each $o \in \hat{O}$ and $\mu(o) \neq \mu(o')$ for $o \neq o'$.

Now consider the base economy. With positive probability, f^1 has a priority ordering, $(\mu(o_1), \dots, \mu(o_m), i^{**})$ followed by an arbitrary permutation of the remaining agents. Given such a priority ordering, the objects in \hat{O} will be all consumed before i^{**} gets her turn but b will not be consumed before i^{**} gets her turn, so she will consume b . This proves that $a \triangleright (RP^1, m^1) b$. \square

Lemma 3. RP^1 is wasteful if and only if RP^q is wasteful for any $q \in \mathbb{N}$.

Proof. We proceed in two steps.

(i) the “only if” Part: Suppose that RP^1 is wasteful. Then, there are objects $a, b \in \tilde{O}$ and an agent $i^* \in N^1$ who prefers a over b such that she consumes b under some ordering $(\tilde{i}_{(1)}^1, \dots, \tilde{i}_{(|N^1|)}^1)$ (implied by some \tilde{f}^1) and that a is not consumed by any agent

under $(\hat{i}_{(1)}^1, \dots, \hat{i}_{(|N^1|)}^1)$ (implied by some \hat{f}^1). (This is the necessary implication of the “wastefulness” under RP^1 .)

Now consider its q -fold replica, RP^q . With positive probability, an ordering $(\bar{\gamma}(\tilde{i}_{(1)}^1), \dots, \bar{\gamma}(\tilde{i}_{(|N^1|)}^1))$ arises, where $\bar{\gamma}(i)$ is an arbitrary permutation of $\gamma(i)$. Clearly, each agent in $\gamma(i^*)$ must consume b even though she prefers a over b (since all copies of all goods the agents in $\gamma(i^*)$ prefers to b are all consumed by the agents ahead of them). Likewise, with positive probability, an ordering $(\bar{\gamma}(\hat{i}_{(1)}^1), \dots, \bar{\gamma}(\hat{i}_{(|N^1|)}^1))$ arises. Clearly, under this ordering, no copies of good a are consumed. It follows that RP^q is wasteful.

(ii) the “if” Part: Suppose next that RP^q is wasteful. Then, there are objects $a, b \in \tilde{O}$ and an agent $i^{**} \in N^q$ who prefers a over b such that she consumes b under some ordering $(\tilde{i}_{(1)}^q, \dots, \tilde{i}_{(|N^q|)}^q)$ (implied by some \tilde{f}^q) and that not all copies of good a are consumed by any agent under $(\hat{i}_{(1)}^q, \dots, \hat{i}_{(|N^q|)}^q)$ (implied by some \hat{f}^q).

Now consider the corresponding base economy and associated RP^1 . The argument of Part (ii) of Lemma 2 implies that there exists an ordering $(\tilde{i}_{(1)}^1, \dots, \tilde{i}_{(|N^1|)}^1)$ under which agent $\tilde{i}^* = \gamma^{-1}(i^{**}) \in N^1$ consumes b even though she prefers a over b .

Next, we prove that RP^1 admits a positive-probability ordering under which good a is not consumed. Let $N'' := \{r \in N^1 \mid \exists j \in \gamma(r) \text{ who consumes the null good under } \hat{f}^q\}$. For each $r \in N''$, we let \emptyset^r denote the null good some clone of $r \in N^1$ consume. In other words, we use different notations for the null good consumed by the clones of different agents in N'' . Given this convention, there can be at most q copies of each \emptyset^r .

Let $\bar{O} := O \cup (\cup_{r \in N''} \emptyset^r) \setminus \{a\}$, and define a correspondence $\psi : N^1 \rightarrow \bar{O}$ by

$$\psi(r) := \{b \in \bar{O} \mid \exists j \in \gamma(r) \text{ who consumes } b \text{ under } \hat{f}^q\}.$$

Claim 11. For each $N' \subset N^1$, $|\cup_{r \in N'} \psi(r)| \geq |N'|$.

Proof. Suppose not. Then, $k := |\cup_{r \in N'} \psi(r)| < |N'| =: l$. Reindex the sets so that $\cup_{r \in N'} \psi(r) =: \{o^1, \dots, o^k\}$ and $N' = \{r^1, \dots, r^l\}$. Let x_{ij} denote the number of copies of object $o^j \in \psi(r^i)$ consumed by the clones of r^i in the q -fold replica under \hat{f}^q .

Since there are at most q copies of each good, we must have

$$\sum_{j=1}^k \sum_{i=1}^l x_{ij} \leq kq.$$

At the same time, all q clones of each agent in N' , excluding $q - 1$ agents (who may be consuming a), are consuming some goods in O' under \hat{f}^q , we must have

$$\sum_{i=1}^l \sum_{j=1}^k x_{ij} \geq lq + q - 1 = (l - 1)q + 1 > kq,$$

We thus have a contradiction. \parallel

Claim 11 then implies, via Hall's theorem, that there exists a mapping $\nu : N^1 \rightarrow \bar{O}$ such that $\nu(r) \in \psi(r)$ for each $r \in N^1$ and $\nu(r) \neq \nu(r')$ if $r \neq r'$.

Let $O' \subset \bar{O}$ be the subset of all object types in \bar{O} whose entire q copies are consumed under \hat{f}^q . Order O' in the order that the last copy of each good is consumed; i.e., label $O' = \{o^1, \dots, o^m\}$ such that the last copy of good o^i is consumed prior to the last copy of o^j if $i < j$. Let \hat{N} be any permutation of the agents in $\nu^{-1}(\bar{O} \setminus O')$. Now consider the ordering in RP^1 : $(\hat{i}_{(1)}^1, \dots, \hat{i}_{(|N^1|)}^1) = (\nu^{-1}(o^1), \dots, \nu^{-1}(o^m), \hat{N})$, where the notational convention is as follows: for any $l \in \{1, \dots, m\}$, if $\nu^{-1}(o^l)$ is empty, then no agent is ordered.

Claim 12. Under the ordering $(\hat{i}_{(1)}^1, \dots, \hat{i}_{(|N^1|)}^1) = (\nu^{-1}(o^1), \dots, \nu^{-1}(o^m), \hat{N})$, a is not consumed.

Proof. For any $l = 0, \dots, m$, let O^l be the set of objects that are consumed by agents $\nu^{-1}(o^1), \dots, \nu^{-1}(o^l)$ under the current ordering (note that some of $\nu^{-1}(o^1), \dots, \nu^{-1}(o^l)$ may be nonexistent). We shall show $O^l \subseteq \{o^1, \dots, o^l\}$ by an inductive argument. First note that the claim is obvious for $l = 0$. Assume that the claim holds for $0, 1, \dots, l-1$. If $\nu^{-1}(o^l) = \emptyset$, then no agent exists to consume a good at this step and hence the claim is obvious. Suppose $\nu^{-1}(o^l) \neq \emptyset$. By definition of ν , agent $\nu^{-1}(o^l)$ weakly prefers o^l to any good in $\bar{O} \setminus \{o^1, \dots, o^{l-1}\}$. Therefore $\nu^{-1}(o^l)$ consumes a good in $\{o^l\} \cup (\{o^1, \dots, o^{l-1}\} \setminus O^{l-1}) \subseteq \{o^1, \dots, o^l\}$. This and the inductive assumption imply $O^l \subseteq \{o^1, \dots, o^l\}$.

Next, consider agents that appears in the ordered set \hat{N} . By an argument similar to the previous paragraph, each agent i in \hat{N} consumes an object in $\nu(i) \cup (\{o^1, \dots, o^m\} \setminus O^m)$. In particular, no agent in \hat{N} consumes a . \parallel

Since the ordering $(\hat{i}_{(1)}^1, \dots, \hat{i}_{(|N^1|)}^1) = (\nu^{-1}(o^1), \dots, \nu^{-1}(o^m), \hat{N})$ realizes with positive probability under RP^1 , Claim 12 completes the proof of Lemma 3. \square

Proof of Proposition 3. If RP^q is ordinally inefficient for some $q \in \mathbb{N}$, then either it is wasteful or there must be a cycle of binary relation $\triangleright(RP^q, m^q)$. Lemmas 2 and 3 then imply that RP^1 is wasteful or there exists a cycle of $\triangleright(RP^1, m^1)$, and that $RP^{q'}$ is wasteful or there exists a cycle of $\triangleright(RP^{q'}, m^{q'})$ for each $q' \in \mathbb{N}$. Hence, for each $q' \in \mathbb{N}$, $RP^{q'}$ is ordinally inefficient. \square

REFERENCES

ABDULKADIROĞLU, A., Y.-K. CHE, AND Y. YASUDA (2008): "Expanding 'Choice' in School Choice," mimeo.

- ABDULKADIROĞLU, A., P. A. PATHAK, AND A. E. ROTH (2005): “The New York City High School Match,” *American Economic Review Papers and Proceedings*, 95, 364–367.
- ABDULKADIROĞLU, A., P. A. PATHAK, A. E. ROTH, AND T. SÖNMEZ (2005): “The Boston Public School Match,” *American Economic Review Papers and Proceedings*, 95, 368–372.
- ABDULKADIROĞLU, A., AND T. SÖNMEZ (1998): “Random Serial Dictatorship and the Core from Random Endowments in House Allocation Problems,” *Econometrica*, 66, 689–701.
- (1999): “House Allocation with Existing Tenants,” *Journal of Economic Theory*, 88, 233–260.
- (2003a): “Ordinal Efficiency and Dominated Sets of Assignments,” *Journal of Economic Theory*, 112, 157–172.
- (2003b): “School Choice: A Mechanism Design Approach,” *American Economic Review*, 93, 729–747.
- BALINSKI, M., AND T. SÖNMEZ (1999): “A tale of two mechanisms: student placement,” *Journal of Economic Theory*, 84, 73–94.
- BOGOMOLNAIA, A., AND H. MOULIN (2001): “A New Solution to the Random Assignment Problem,” *Journal of Economic Theory*, 100, 295–328.
- (2002): “A Simple Random Assignment Problem with a Unique Solution,” *Economic Theory*, 19, 623–635.
- CHEN, Y., AND T. SÖNMEZ (2002): “Improving Efficiency of On-campus Housing: An Experimental Study,” *American Economic Review*, 92, 1669–1686.
- CRÈS, H., AND H. MOULIN (2001): “Scheduling with Opting Out: Improving upon Random Priority,” *Operations Research*, 49, 565–577.
- CRIPPS, M., AND J. SWINKELS (2006): “Efficiency of Large Double Auctions,” *Econometrica*, 74, 47–92.
- GALE, D., AND L. S. SHAPLEY (1962): “College Admissions and the Stability of Marriage,” *American Mathematical Monthly*, 69, 9–15.
- GRESIK, T., AND M. SATTERTHWAITTE (1989): “The Rate at Which a Simple Market Converges to Efficiency as the Number of Traders Increases,” *Journal of Economic Theory*, 48, 304–332.
- HURWICZ, L. (1972): “On Informationally Decentralized Systems,” in *Decision and Organization*, ed. by G. McGuire, and R. Radner. North Holland, Amsterdam.
- HYLLAND, A., AND R. ZECKHAUSER (1979): “The Efficient Allocation of Individuals to Positions,” *Journal of Political Economy*, 87, 293–314.
- IMMORLICA, N., AND M. MAHDIAN (2005): “Marriage, Honesty, and Stability,” *SODA 2005*, pp. 53–62.
- JACKSON, M. O. (1992): “Incentive compatibility and competitive allocations,” *Economics Letters*, pp. 299–302.
- JACKSON, M. O., AND A. M. MANELLI (1997): “Approximately competitive equilibria in large finite economies,” *Journal of Economic Theory*, pp. 354–376.
- JUDD, K. L. (1985): “The Law of Large Numbers with a Continuum of IID Random Variables,” *Journal of Economic Theory*, 35, 19–25.
- KATTA, A.-K., AND J. SETHURAMAN (2006): “A Solution to The Random Assignment Problem on The Full Preference Domain,” forthcoming, *Journal of Economic Theory*.
- KESTEN, O. (2008): “Why do popular mechanisms lack efficiency in random environments?,” Carnegie Mellon University, Unpublished mimeo.

- KOJIMA, F. (2008): “Random Assignment of Multiple Indivisible Objects,” forthcoming, *Mathematical Social Sciences*.
- KOJIMA, F., AND M. MANEA (2008): “Strategy-proofness of the Probabilistic Serial Mechanism,” Harvard University, Unpublished mimeo.
- KOJIMA, F., AND P. A. PATHAK (2008): “Incentives and Stability in Large Two-Sided Matching Markets,” forthcoming, *American Economic Review*.
- MANEA, M. (2006a): “Asymptotic Ordinal Inefficiency of Random Serial Dictatorship,” Harvard University, Unpublished mimeo.
- (2006b): “A Constructive Proof of The Ordinal Efficiency Welfare Theorem,” forthcoming, *Journal of Economic Theory*.
- MCLENNAN, A. (2002): “Ordinal Efficiency and The Polyhedral Separating Hyperplane Theorem,” *Journal of Economic Theory*, 105, 435–449.
- PATHAK, P. A. (2006): “Lotteries in Student Assignment,” Harvard University, unpublished mimeo.
- ROBERTS, D. J., AND A. POSTLEWAITE (1976): “The Incentives for Price-Taking Behavior in Large Exchange Economies,” *Econometrica*, 44, 115–127.
- ROTH, A. E., AND E. PERANSON (1999): “The Redesign of the Matching Market for American Physicians: Some Engineering Aspects of Economic Design,” *American Economic Review*, 89, 748–780.
- RUSTICHINI, A., M. SATTERTHWAITE, AND S. WILLIAMS (1994): “Convergence to Efficiency in a Simple Market with Incomplete Information,” *Econometrica*, 62, 1041–1064.
- SUN, Y. (2006): “The Exact Law of Large Numbers via Fubini Extension and Characterization of Insurable Risks,” *Journal of Economic Theory*, 126, 31–69.
- YILMAZ, O. (2006): “House Allocation with Existing Tenants: a New Solution,” University of Rochester, Unpublished mimeo.